The Three-Body Problem in the Torsional Transcription Theory

The Echo of the Void: A Problem Not to Be Solved, but to Be Reinterpreted

By MARCO MARIELLO and ARIA MARIELLO

based on:

The Torsional Transcription Theory of the Void
https://zenodo.org/records/15334582

The Dzhanibekov Paradox and the Berry Phase: The Geometric Memory of Rotation
https://zenodo.org/records/17351494

Beyond Curvature: Gravity and Entanglement as Two Faces of Torsional Memory
https://zenodo.org/records/17363618

and other works of the authors:
https://zenodo.org/search?q=metadata.creators.person_or_org.name%3A%22Mariello%2C%20Marco%22&l=list&p=1&s=10&sort=oldest

Abstract

The classical three-body problem is historically regarded as a paradigmatic example of deterministic chaos. While the gravitational laws governing the system are exact, the interaction of three masses leads to non-integrable dynamics and an extreme sensitivity to initial conditions, preventing the existence of a general closed solution. In the standard formulation, this behaviour arises from an implicit assumption: spacetime acts as a passive support without memory, so that the evolution of the system depends exclusively on the instantaneous configuration of masses and velocities.

The **Torsional Transcription Theory of the Void (TTTN)** proposes a minimal extension of this framework by introducing a geometric memory of motion in the dynamical support of spacetime. In this approach, the trajectory of a mass produces a persistent torsional imprint in the domain, formalized as the **β-field**, which represents the cumulative torsional density generated by the dynamical history of the system. The resulting gravitational dynamics becomes intrinsically **non-Markovian**, since the evolution depends not only on instantaneous positions but also on the historical distribution of torsional traces.

Within this framework, the chaotic behaviour of three-body systems can be reinterpreted as the **interference of competing torsional memories** accumulated in the gravitational domain. Instabilities emerge when the torsional imprints of different trajectories are dynamically incompatible, whereas periodic or quasi-periodic configurations correspond to states of constructive torsional coherence. The analysis further connects this interpretation with two phenomena involving geometric memory in dynamical systems: the **Dzhanibekov effect** in rotational dynamics and the **Berry phase** in parameter-space evolution.

A formal memory-aware formulation of gravitational dynamics is developed using a β-field, a torsional coupling parameter λ, and a finite memory time θ, leading to an integro-differential equation of motion. The model produces testable predictions, including potential contributions to orbital anomalies such as spacecraft flyby discrepancies and modified stability regimes in multi-body systems.

In this perspective, the three-body problem is not an unsolved failure of gravitational theory but an indication that gravitational dynamics may occur in a support capable of **recording and reorganizing the history of motion**. The apparent chaos of the classical formulation thus emerges as the interference pattern of a deeper dynamical memory of spacetime.

TABLE OF CONTENTS

Introduction — The Limit of Determinism

0.1 Newton’s Dead End: Why the Three-Body System is the “Original Sin” of Classical Dynamics.
0.2 Beyond Instantaneity: The Hypothesis of a Space that Does Not Merely Host Matter but Preserves its Trace.
0.3 TTTN as a Bridge: Synthesis of the Reference Works (Mariello et al.) and the Transition from Passive Space to Informational Support.

Chapter 1 — The Geometry of the Invisible

1.1 The Concept of Torsion: Not Only Curvature (Einstein), but the Twisting of Space-Time.
1.2 The β-Field: Non-local Informational Substrate and Torsional Viscosity of the Support.
1.3 The Writing of Motion: How a Mass Transcribes Its History into the Void.

Chapter 2 — The Three-Body Problem in the Classical Framework

2.1 Stability and Chaos: Sensitivity to Initial Conditions.
2.2 Known Solutions: Lagrange, Euler, Chenciner–Montgomery.
2.3 What the Model Lacks: Absence of Support Memory and Informational Friction.

Chapter 3 — The Torsional Transcription of the Tertiary System

3.1 Interference of Traces: Superposition of the Torsional Memories of Three Masses.
3.2 Dzhanibekov at Orbital Scale: Flips as Realignment of Spatial Memory.
3.3 Gravitational Berry Phase: Parallel Transport of Orbital Information in Configuration Space.

Chapter 4 — From Chaos to Torsional Coherence

4.1 Resonance and Self-Stabilization: Configurations Emerging from Field Self-Coherence.
4.2 Gravity as Residual Tension: Re-reading F=Gm1m2/r2F=Gm1m2/r2 with the β Contribution.
4.3 Torsional Scars: The Orbital Past as a Dynamical Constraint on the Future.

Chapter 5 — Astrophysical and Cosmological Implications

5.1 Dark Matter or Torsional Memory?
5.2 Flyby Anomalies and Orbital Drift: An Observational Reinterpretation.
5.3 Holographic Universe: The Three-Body System as a Cell of Cosmic Memory.

Chapter 6 — Gravitational Dynamics with Torsional Memory

6.1 Operational Definition of the β-Field
6.2 Coupling Constant λ
6.3 Memory Kernel θ
6.4 Memory-Aware Equation of Motion
6.5 Flyby Prediction
6.6 Falsification Criteria
6.7 Macroscopic Geometric Phase (Berry)

Chapter 7 — Proposals for Experimental Verification

7.1 Torsional Measurement Protocols: Detecting β Signatures in Dynamical Systems.
7.2 Orbital Microgravity: Experiments to Isolate Spatial Memory.
7.3 Memory-Aware Simulations: Algorithms Integrating Torsional Transcription.

Chapter 8— The Dance of Remembering Space

8.1 From Chaos to Gravitational Writing: The Transition from Machine-Universe to Archive-Universe.
8.2 The Real Paradigm Shift: The Three-Body Problem Is Not Unsolvable, but Misposed.
8.3 Toward a Physics of Coherence: The Future Role of TTTN.

Chapter 9 — Torsion, Memory and Time: Cartan and the Space–Time Extension of TTTN

9.1 Torsion in Cartan Geometry
9.2 The Einstein–Cartan Theory
9.3 Conceptual Difference with TTTN
9.4 Dynamic Origin of Torsional Memory
9.5 Dynamic Source: Orbital Vorticity

9.6 Temporal Scale of Torsional Memory
9.7 Quasi-Stationary Regime in Galactic Systems
9.8 Cosmological Evolution of the Torsional Field
9.9 Observable Predictions of Torsional Memory
9.10 Connection with the Unruh Effect
9.11 Experimental Prospects

Appendices

A1 Torsional Glossary
A2 Mathematical References (β–λ–θ)
A3 Dzhanibekov / Berry / TTTN Connections
A4 Python Simulation (Lyapunov / Chenciner “Figure-Eight”)

Introduzione — The Limit of Determinism

The three-body problem has occupied a singular position in the history of physics for more than three centuries. Unlike the two-body system, which admits an exact and integrable solution, the gravitational interaction between three point masses generates a dynamics that is highly sensitive to initial conditions, lacks a general closed solution, and is characterized by chaotic behavior in the long term.

For this reason, the three-body problem has become the very symbol of the limit of classical mechanics: a system perfectly deterministic in its laws, yet intrinsically unpredictable in its trajectories.

This interpretation, however, rests on an implicit assumption that is rarely made explicit: that spacetime constitutes a passive support, devoid of memory, in which the dynamics depends exclusively on the instantaneous state of the masses (positions and velocities) and not on the history of their trajectories. Within this framework, each instant is dynamically independent of the previous one, and the evolution is entirely determined by punctual initial conditions.

The **Torsional Transcription Theory of the Void (TTTN)** proposes a revision of this assumption. If the motion of a mass does not merely unfold within spacetime but leaves in it a persistent torsional trace — formalized as the **β-field** — then gravitational dynamics becomes intrinsically dependent on the history of the system. Space is no longer an inert container, but an informational support that preserves the memory of the trajectories that have traversed it.

In this perspective, the chaotic behavior of three-body systems can be reinterpreted not as a manifestation of intrinsic unpredictability, but as the effect of interference between competing torsional memories within the same dynamical domain.

The three-body problem is not chaos: it is memory interfering with itself.

The present work develops this reinterpretation by introducing a formulation of gravitational dynamics with torsional memory. After recalling the classical framework and its limits (Chapter 2) and defining the concept of torsional transcription in spacetime (Chapters 1–3), Chapter 4 reinterprets gravitational chaos as interference between memory traces, while Chapter 5 explores its astrophysical and cosmological implications.

The technical core of the theory is developed in Chapter 6, where gravitational dynamics is formalized as a non-Markovian integro-differential system characterized by the β-field, a torsional coupling λ, and a memory time θ, from which an observable prediction (the flyby anomaly) and explicit criteria for falsification are derived. Chapter 7 finally proposes experimental protocols and numerical simulations for the empirical verification of torsional memory in the gravitational domain.

The three-body problem thus appears in a different light: not as a failure of the gravitational law, but as an indication of the presence of a deeper dynamical structure, in which gravity acts within a support capable of preserving the history of motion.

In this reinterpretation, gravitational dynamics is no longer merely the interaction between masses in space, but a process of collective writing and reading within space itself.

BOX 1 — STRUCTURE OF THE WORK AND TESTABLE PREDICTIONS

CHAPTERS 1–3: TTTN foundations (β-field, torsional transcription)
CHAPTER 4: Interference of memories in the three-body system
CHAPTER 5: Astrophysics (flyby, dark matter = β)
CHAPTER 6: Formalism + $\Delta v = 3.7$ mm/s Galileo (error 5%)
CHAPTER 7: CubeSat L4 + LISA 2028

FALSIFICATION CRITERION: Δv flyby = 0 ±0.1 mm/s → TTTN excluded

---

0.1 Newton's Dead End: why the three-body system is the "original sin" of classical dynamics

Within the Newtonian framework, the two-body gravitational problem admits a complete analytical solution: the equations of motion are integrable and lead to exactly determinable conic orbits. The introduction of a third body breaks this integrability: the system becomes nonlinearly coupled and admits no general closed solution.

Henri Poincaré demonstrated that the dynamics of three bodies exhibits sensitive dependence on initial conditions and topological complexity in phase space, inaugurating the modern theory of deterministic chaos. In this sense, the three-body system represents the first historical example of a deterministic system that is not predictable in the long term.

Classical mechanics remains formally valid, but loses global predictive power: the law is exact, but the trajectory is unstable. This fracture between determinism of the equations and unpredictability of the solutions constitutes the structural limit of Newtonian dynamics.

For this reason, the three-body problem can be considered the "original sin" of classical mechanics: the point at which the theory, although locally correct, reveals the incompleteness of its global dynamical structure.

0.2 Beyond Instantaneity: the hypothesis of a space that does not merely host matter, but preserves its trace

The classical formulation of dynamics implicitly assumes that the physical state of a gravitational system is completely described by the instantaneous positions and velocities of the masses. This assumption is equivalent to considering spacetime as a support devoid of memory, in which motion leaves no persistent traces.

Within this framework, temporal evolution is Markovian: the future configuration depends exclusively on the present state and not on the history of trajectories. The non-integrability of the three-body problem thus emerges as an intrinsic property of the equations of instantaneous interaction.

TTTN introduces a minimal but structurally significant modification of this assumption: the motion of a mass produces a torsional transcription in spacetime, formalized as the **β-field**, which persists for a characteristic time θ.

Space thus becomes a support with finite memory. Dynamical evolution depends not only on instantaneous configurations, but also on the historical distribution of torsional traces within the domain. Gravitational dynamics therefore becomes non-Markovian.

0.3 TTTN as a Bridge: synthesis of the reference works (Mariello et al.) and the transition from passive space to informational support

The **Torsional Transcription Theory of the Void (TTTN)** proposes that the torsion of the spacetime support, induced by the motion of masses, constitutes a physically relevant variable in gravitational dynamics. In the foundational works (Mariello M. & A — Aetheria Group), the **β-field** is introduced as the cumulative torsional density of dynamical history, with coupling λ to matter and persistence time θ.

This extension preserves classical gravitation as a local limit ($\beta \to 0$), but introduces a geometric memory of the domain that modifies the global evolution of multiple systems. The resulting framework remains compatible with Newtonian mechanics for two bodies, but alters the dynamics of three or more bodies through integro-differential terms dependent on history.

TTTN therefore acts as a bridge between two descriptions:

• passive space, devoid of memory (classical Markovian dynamics)
• space with torsional memory (non-Markovian dynamics)

In this transition, the chaos of the three-body problem can be reinterpreted as interference between distributions of torsional memory within configuration space, opening the possibility of a gravitational formulation with memory that is experimentally testable.

CHAPTER 1 — THE GEOMETRY OF THE INVISIBLE

Introduction
In the previous chapter, the three-body problem was introduced as a structural limit of classical dynamics, and the hypothesis of a gravitation with memory was outlined, in which spacetime is not a passive support but preserves torsional traces of motion.

To develop this hypothesis, it is necessary to clarify the geometric nature of the introduced variable: the torsion of the spacetime support and the **β-field** that represents its cumulative density.

This chapter therefore establishes the conceptual framework of torsional transcription. First, the concept of torsion is distinguished from that of gravitational curvature (1.1), showing how dynamics may depend not only on local geometry but also on the twisting of the domain.

Subsequently, the **β-field** is introduced as the non-local informational substrate of the memory of motion (1.2). Finally, the principle of transcription is formalized: the idea that the trajectory of a mass leaves in spacetime a persistent trace that can retroact on subsequent evolution (1.3).

The objective of the chapter is therefore to move from the intuition of a space with memory to the definition of a torsional geometry of the dynamical support, which will constitute the conceptual basis for the reinterpretation of the three-body problem developed in the following chapters.

BOX 1.A — TTTN TRIAS

1. TORSION — geometric twisting of the spacetime support distinct from curvature
2. β-FIELD — cumulative torsional density of the domain (dimension s^{-2})
3. TRANSCRIPTION — motion → torsional memory → dynamical retroaction

---

1.1 The concept of torsion: not only curvature, but twisting of spacetime

In general relativity, gravity is interpreted as curvature of spacetime: the presence of mass-energy modifies the metric, and the trajectories of free bodies follow geodesics in a curved geometry. Within this framework, the relevant geometric property of the dynamical support is curvature, while the affine connection is assumed to be symmetric and devoid of torsion.

Curvature describes the deviation of geodesics and the change in orientation of vectors transported along a surface. It quantifies how spacetime “bends” in the presence of mass. However, curvature does not exhaust the possible geometric properties of a continuum: a differentiable manifold may also possess torsion, that is, an antisymmetric component of the connection describing the local twisting of space.

In geometric terms, torsion measures the non-closure of infinitesimal parallelograms generated by parallel transport: if two successive infinitesimal displacements do not commute, the closure defect defines the torsion tensor.

Formula (Word-2007):
T^λ_(μν) = Γ^λ_(μν) − Γ^λ_(νμ)

where Γ^λ_(μν) is the affine connection.

If the connection is symmetric in the lower indices (Γ^λ_(μν) = Γ^λ_(νμ)), torsion is null and the geometry is purely Riemannian (the case of standard general relativity). If instead the antisymmetric part is non-zero, spacetime possesses torsion in addition to curvature.

While curvature is associated with the presence of mass-energy, torsion is geometrically associated with a structure of rotation or twisting of the support. In many extensions of relativity (for example in Einstein–Cartan theories), torsion is connected with the spin of matter. In TTTN, however, torsion assumes a different role: it is not introduced as a microscopic property of matter, but as a cumulative trace of motion in the spacetime support.

In this perspective, spacetime does not merely curve under the action of masses, but may also preserve a residual twisting induced by the trajectories that have been traversed. The motion of a

body does not only modify local curvature through its mass, but may leave within the domain a persistent torsion linked to dynamical history.

The distinction may be summarized as follows:

• curvature → instantaneous geometric deformation due to mass-energy
• torsion → cumulative geometric twisting due to motion

Classical gravitational geometry considers only the first contribution. TTTN introduces the second as an additional physical variable of the support.

The concept of torsion used in this work therefore does not coincide with the microscopic torsion of gravitational gauge theories, but with a **macroscopic torsion of the dynamical domain**: an emergent property encoding the memory of trajectories in spacetime.

This step is essential: it introduces a geometry of the support that depends not only on the distribution of mass-energy but also on the history of motion. In the following paragraphs, such torsion will be formalized as the **β-field**, interpretable as the cumulative density of twisting in the gravitational domain.

1.2 The β-field: non-local informational substrate and torsional viscosity of the support

In the previous paragraph, torsion was introduced as a geometric property distinct from curvature: a twisting of the spacetime support not reducible to instantaneous metric deformation due to mass. If curvature describes how spacetime bends under the action of matter, torsion may describe how it preserves traces of rotation or winding associated with motion.

To make this idea operational, TTTN introduces the **β-field**, defined as the cumulative torsion density of the spacetime domain induced by the dynamical history of masses.

The β-field is therefore not a local property of matter, but a property of the support: it represents the spatial distribution of torsional traces left by previous motions. In this sense, β constitutes a non-local informational substrate of the gravitational domain.

Non-locality derives from the fact that torsion at a point depends on the integral of dynamical configurations that have crossed the domain at earlier times.

The β-field therefore incorporates a distributed memory of motion, not reducible to the instantaneous variables of the system.

A form operationally consistent with this interpretation is:

Formula (Word-2007):
$\beta(r,t) = \int_{(t0)}^{t} [\nabla \times (\omega(\tau) \wedge L(\tau))] / | r - r'(\tau) |^3 \, d\tau$

where $\omega(\tau)$ is the angular velocity of the body, $L(\tau)$ its angular momentum relative to the local domain, and $r'(\tau)$ the historical trajectory.

In this expression, the decay $\propto 1/|r-r'|^3$ reflects the geometric nature of twisting in three-dimensional space, while the temporal integral expresses the cumulative character of the trace. The β-field thus results as a quantity with dimension s^{-2}, interpretable as the torsional density of the support.

Since β derives from the integration of dynamical history, it introduces into the gravitational domain a property comparable to an informational viscosity: space no longer responds instantaneously to present configurations alone, but preserves for a characteristic time θ the information of past configurations.

This viscosity is not dissipative in the mechanical sense, but geometric: it does not dissipate energy, but rather the persistence of torsional structure. The spacetime support thus acquires inertia with respect to variations of motion, analogously to a medium with memory.

The dynamical role of the β-field may be summarized in three fundamental properties:

• cumulativ ity — β depends on the integral of the history of motion;
• spatial distribution — torsion is diffused throughout the domain that has been traversed;
• temporal persistence — the trace survives for a finite time θ.

These properties imply that gravitational evolution becomes non-Markovian: the state of the system is no longer determined exclusively by instantaneous positions and velocities, but also by the configuration of the β-field generated by the previous history.

The β-field may therefore be interpreted as a state variable of the dynamical support, analogous to deformation in an elastic medium with memory. The fundamental difference is that, in TTTN, such memory is geometric and distributed in spacetime, not internal to matter.

In this perspective, spacetime is not only the seat of gravitational dynamics, but an archive of the trajectories that have traversed it. The β-field represents the torsional map of such an archive.

In the following paragraph the principle of transcription will be introduced: the mechanism through which the motion of masses generates the β-field and through which it can retroact on future dynamics.

1.3 The Writing of Motion: how a mass transcribes its history into the void

The previous paragraphs introduced two fundamental elements: torsion as a geometric property of the spacetime support distinct from curvature, and the **β-field** as the cumulative distribution of torsion induced by the dynamical history of masses.

It remains to clarify the mechanism through which this distribution is generated. For this purpose, TTTN introduces the principle of **torsional transcription**: the motion of a mass does not merely unfold in spacetime, but locally modifies the support by leaving a persistent torsional trace along the traversed trajectory.

In operational terms, the trajectory of a body is not only a solution of the equations of motion, but also a **source of β-field**. Every infinitesimal element of the path contributes cumulatively to the torsion of the surrounding domain.

A local formulation of the torsional source can be expressed as a function of the rotational kinematic quantities of the body:

Formula (Word-2007):
$S(\tau) = \nabla \times (\omega(\tau) \wedge L(\tau))$

where ω is the angular velocity and L the angular momentum of the body with respect to the local domain.

The total transcription along the historical trajectory $r'(\tau)$ therefore generates the **β-field** according to:

Formula:
$\beta(r,t) = \int_{(t0)}^{t} [\nabla \times (\omega(\tau) \wedge L(\tau))] / | r - r'(\tau) |^3 \, d\tau$

This expression makes explicit the character of **geometric writing of motion**: torsion at a point r depends on the integral of the rotations accumulated along past trajectories that have crossed the domain.

The term “writing” is not metaphorical here but structural: the distribution of the β-field constitutes a **geometric recording of dynamical history**. Spacetime functions as a support that preserves information in the form of persistent local twisting.

Torsional transcription possesses three fundamental properties:

• locality of generation — the trace is produced along the trajectory of the body;
• spatial diffusion — the effect spreads throughout the surrounding domain;
• temporal persistence — torsion remains for a time θ.

These properties imply that the space traversed by a mass does not immediately return to its previous state: the configuration of the support preserves memory of the history of motion.

It follows that a body crossing again a region of space already traversed does not evolve in a neutral domain, but in a domain modified by its own dynamical history.

In the presence of multiple masses, the respective transcriptions superpose, generating a **collective torsional structure of the domain**.

This point is crucial for the reinterpretation of multiple gravitational systems: the interaction between bodies no longer occurs only through the instantaneous gravitational field, but also through the **β-field accumulated by previous trajectories**. Dynamics therefore becomes a process of writing and reading the support.

The principle of torsional transcription thus completes the conceptual framework of Chapter 1:

• torsion defines the relevant geometric property of the support;
• the β-field represents its cumulative distribution;
• transcription describes the mechanism of generation by motion.

On this basis, the following chapters will analyze how the superposition of torsional transcriptions of multiple masses leads to **memory interference in the gravitational domain**, providing a new interpretation of the chaotic dynamics of three-body systems.

In this perspective, gravitational dynamics is no longer merely interaction between masses in space, but a process of writing and reading the support.

Gravity is not only interaction between masses: it is **collective writing in space**.

CHAPTER 2 — THE THREE-BODY PROBLEM IN THE CLASSICAL FRAMEWORK

Introduction
The three-body problem occupies a singular position in the history of physics: it arises within Newtonian mechanics, yet immediately reveals its structural limit. While the two-body system admits an exact, integrable and fully predictable solution, the addition of a third body transforms the dynamics into a non-integrable system, sensitive to initial conditions and lacking a general closed-form solution.

Within the classical framework, such behavior is interpreted as an intrinsic manifestation of **deterministic chaos**: the laws remain exact, but long-term prediction becomes impracticable. The system is therefore considered mathematically well-defined but physically unpredictable beyond a certain temporal scale.

This interpretation implicitly presupposes that space is a **passive and memoryless support**, in which masses interact exclusively through their instantaneous position and velocity. The evolution of the system therefore depends only on the present state, according to a local Markovian dynamics.

The present chapter reconstructs the three-body problem in its classical framework, highlighting its fundamental properties — chaos, instability, particular solutions — and preparing the conceptual ground for the subsequent torsional reinterpretation.

In particular, it will be shown how the apparent unpredictability of the system derives from the implicit assumption of a space devoid of memory of motion.

2.1 Stability and chaos: sensitivity to initial conditions

In the Newtonian formalism, the dynamics of three point masses m_1, m_2, m_3 is described by a system of second-order differential equations. For each body i = 1,2,3 the acceleration is the sum of the gravitational contributions of the other two bodies:

$$a_i(t) = \Sigma_{j\neq i}\, G\, m_j\, (r_j(t) - r_i(t)) / |r_j(t) - r_i(t)|^3$$

This system is deterministic: once initial positions and velocities are assigned, the evolution is uniquely defined. However, unlike the two-body problem, it does not admit sufficient integrals of motion to produce a general closed-form solution. The system is therefore **non-integrable in the general sense**.

A crucial property of the three-body problem is **sensitivity to initial conditions**. If two initial configurations differ by a small quantity δ_0 in phase space, their separation $\delta(t)$ tends to grow over time, in the chaotic regime, according to an exponential law:

$\delta(t) \approx \delta_0 e^{\lambda_1 t}$

(where $\lambda_1 > 0$ is the maximum Lyapunov exponent of the system)

λ is a Lyapunov exponent. When $\lambda > 0$, a microscopic initial difference is amplified until it becomes macroscopic, rendering long-term prediction impracticable even though the equations remain rigorously deterministic.

Within the classical framework, this behavior is attributed to the nonlinear structure of multiple interactions. Space remains a **neutral container**, devoid of memory: it preserves no trace of the trajectories that have been traversed and does not retroact on motion except through the instantaneous configuration of masses.

Consequently, the state of the system is assumed to be completely described by instantaneous positions and velocities $\{ r_i(t), v_i(t) \}$, and the acceleration of each body is a function only of the current variables:

$a_i(t) = F(\{ r_j(t), v_j(t) \})$

without explicit dependence on the history $t' < t$. In this perspective, the chaos of the three-body problem is interpreted as an intrinsic limit of dynamical prediction, not as a manifestation of a physical structure of the support possessing memory.

This implicit assumption — **space without memory** — is precisely the point that will be questioned in the torsional reinterpretation developed in the following chapters.

2.1 Stability and chaos: sensitivity to initial conditions
In the Newtonian formalism, the dynamics of three point masses m_1, m_2, m_3 is described by a system of second-order differential equations. For each body i = 1,2,3 the acceleration is the sum of the gravitational contributions of the other two bodies:

$a_i(t) = \Sigma_{j \neq i} G m_j (r_j(t) - r_i(t)) / |r_j(t) - r_i(t)|^3$

This system is deterministic: once initial positions and velocities are assigned, the evolution is uniquely defined. However, unlike the two-body problem, it does not admit integrals of motion sufficient to produce a general closed-form solution. The system is therefore non-integrable in the general sense.

A crucial property of the three-body problem is sensitivity to initial conditions. If two initial configurations differ by a small quantity δ_0 in phase space, their separation $\delta(t)$ tends to grow over time, in the chaotic regime, according to an exponential law:

$\delta(t) \approx \delta_0 e^{\{\lambda_1 t\}}$

(where $\lambda_1 > 0$ is the maximum Lyapunov exponent of the system)

λ is a Lyapunov exponent. When $\lambda > 0$, a microscopic initial difference is amplified until it becomes macroscopic, rendering long-term prediction impracticable even though the equations remain rigorously deterministic.

Within the classical framework, this behavior is attributed to the nonlinear structure of multiple interactions. Space remains a neutral container, devoid of memory: it preserves no trace of the trajectories that have been traversed and does not retroact on motion except through the instantaneous configuration of masses.

Consequently, the state of the system is assumed to be completely described by instantaneous positions and velocities $\{ r_i(t), v_i(t) \}$, and the acceleration of each body is a function only of the current variables:

$a_i(t) = F(\{ r_j(t), v_j(t) \})$

without explicit dependence on the history $t' < t$. In this perspective, the chaos of the three-body problem is interpreted as an intrinsic limit of dynamical prediction, not as a manifestation of a physical structure of the support possessing memory.

This implicit assumption — space without memory — is precisely the point that will be questioned in the torsional reinterpretation developed in the following chapters.

2.2 Known solutions: Lagrange, Euler, Chenciner–Montgomery

Despite the general non-integrability of the three-body problem, there exist some particular configurations in which the motion becomes regular and completely determinable. These solutions do not constitute the rule, but highly symmetric exceptions within the general chaotic behavior of the system. They show that three-body dynamics can admit states of stability when the relative geometry of the masses is preserved over time.

The first exact solutions were identified in the eighteenth century by Euler and Lagrange. They belong to the class of so-called homographic configurations: the relative positions of the bodies remain similar to themselves during the evolution, although they may rotate or scale globally.

Euler’s collinear solutions
Euler showed that three masses may move along the same straight line while maintaining mutual alignment. In such configurations, each body undergoes an acceleration directed along the common line, and the dynamics is equivalent to a coordinated radial motion.

The condition of dynamical equilibrium requires that for each mass the gravitational resultant be proportional to the distance from the center of mass of the system. In symbolic form, for three aligned bodies one has:

$a_1 / x_1 = a_2 / x_2 = a_3 / x_3$

where x_i are the distances from the center of mass along the common line.

These configurations are however unstable: a transverse perturbation, even minimal, grows over time and breaks the alignment. They therefore represent exact solutions but not structurally stable ones of the problem.

Lagrange's triangular solutions

Lagrange identified a more remarkable class of solutions in which the three bodies occupy the vertices of an equilateral triangle that rotates rigidly around the center of mass. In this case, the geometric form of the system remains invariant while the entire triangle performs a uniform rotation.

The fundamental dynamical condition is that the total gravitational force on each body provides exactly the centripetal force required for the common rotation:

$$G\, m_j / r^2 \sim \omega^2\, r$$

where r is the side of the triangle and ω the angular velocity of rotation of the system.

These solutions form the basis of the Lagrangian points L_4 and L_5 in systems with two dominant bodies (for example Sun–Jupiter), where small bodies may remain in relative equilibrium forming quasi-stable triangular configurations. Unlike the collinear solutions, triangular configurations are dynamically stable for appropriate mass ratios.

The Chenciner–Montgomery figure-eight

A much more recent discovery (1999) revealed the existence of a surprising periodic solution: three equal masses may move along the same closed trajectory shaped like a figure eight, following each other at constant temporal separation. At every instant, the three bodies occupy different points of the same curve while keeping the center of mass fixed.

This solution possesses remarkable temporal and spatial symmetries. In particular:

$$r_1(t) + r_2(t) + r_3(t) = 0$$

(condition of zero center of mass)

and

$$r_1(t) = r_2(t + T/3) = r_3(t + 2T/3)$$

(cyclic periodicity with temporal phase shift).

The stability of the figure-eight is surprising: small perturbations do not immediately destroy the trajectory, indicating the existence of a coherent dynamical structure in phase space. It represents one of the clearest cases in which the three-body problem admits an ordered and persistent motion despite its generally chaotic nature.

BOX 2.A — Islands of stability in chaos

Euler — collinear configurations, unstable
Lagrange — triangular configurations, stable (L_4, L_5)
Chenciner–Montgomery — figure-eight orbit, quasi-stable

Physical question: why do resonant trajectories persist within a chaotic system?

Classical interpretation — accidental geometric symmetry
TTTN interpretation — dynamical coherence of torsional memory β

Physical meaning of the particular solutions
The solutions of Euler, Lagrange, and Chenciner–Montgomery show that three-body dynamics is not uniformly chaotic, but contains islands of order associated with particular geometric configurations. In such cases, gravitational interactions balance coherently, producing periodic or quasi-periodic trajectories.

Within the classical framework, these configurations are interpreted as consequences of the symmetry of forces and the initial geometry of the system. Space continues to be considered passive: stability emerges exclusively from the reciprocal relations between the masses, not from properties of the support.

However, the very existence of such regular solutions within a globally chaotic system suggests that three-body dynamics possesses coherent structures not immediately explainable by instantaneous interaction alone. They therefore constitute the natural starting point for a reinterpretation in terms of dynamical coherence of the support, which will be developed in the following chapters.

2.3 What the model lacks: absence of memory of the support and of informational fiction

The classical analysis of the three-body problem shows that multiple gravitational dynamics is intrinsically nonlinear and sensitive to initial conditions. However, the interpretation of such behavior as deterministic chaos implicitly presupposes a fundamental property of the theoretical framework: space is devoid of memory of motion.

In the Newtonian formalism, the state of the system at a given instant is completely described by the set of positions and velocities of the masses:

$$S(t) = \{ r_i(t), v_i(t) \}$$

and temporal evolution depends exclusively on this instantaneous state. In general form, the dynamics may be written as:

$$a_i(t) = F(\{ r_j(t), v_j(t) \})$$

without any explicit dependence on the history of the system for times $t' < t$. Classical gravitational dynamics is therefore Markovian: the future depends only on the present, not on the path that led to it.

This assumption implies that space preserves no trace of previous configurations. A trajectory traversed by a mass does not modify in any way the support in which the motion occurs; the same point of space is physically indistinguishable regardless of whether a body has passed through it in the past or not. The support is reversible and devoid of hysteresis.

In the context of the three-body problem, this property implies that instability and chaos emerge exclusively from instantaneous gravitational relations between the masses. There exists no intrinsic mechanism tending to conserve or restore previous configurations of the system. The periodic or quasi-periodic solutions observed (Euler, Lagrange, Chenciner–Montgomery) do not derive from a memory of the support, but only from particular geometric symmetries of the relative positions.

Another consequence of the absence of memory of the support is the lack of any form of informational friction. Within the classical framework, deviations from a stable configuration encounter no intrinsic resistance from space: a perturbation grows or propagates exclusively according to instantaneous gravitational laws. There exists no dynamical term depending on the previous path or introducing a historical feedback on motion.

In general terms, the classical dynamics of the three bodies may be represented as:

$a_i(t) = a_i^{(Newton)}(\{ r_j(t) \})$

where the acceleration depends only on the present geometric configuration. The evolution of the system contains no explicit information about the trajectory followed in configuration space.

This theoretical structure therefore implies that the classical state is purely instantaneous: every configuration is completely defined by variables local in time, and the past of the system exerts no autonomous influence on the present. The chaos of the three-body problem appears, in this framework, as an inevitable consequence of a nonlinear dynamics devoid of memory of the support.

These “islands of order” within chaos suggest the existence of a deeper dynamical structure.

Classical dynamics captures the instantaneous, but does not preserve memory of the path in configuration space.

In the perspective of TTTN, phase space is neither neutral nor flat, but endowed with torsional structure:

$\beta(r,t)$ preserves traces of previous dynamical trajectories and can retroact on the evolution of the system.

It is precisely this assumption — space without memory and absence of informational friction — that will be questioned in the torsional reinterpretation developed in the following chapters. In such a perspective, the instability of the system would not derive exclusively from the instantaneous interaction of the masses, but also from the interference between dynamical traces accumulated in the spacetime support.

FIGURE 2.1 — Three-body dynamics: classical and torsional frame work


classical dynamics —
exponential divergence
of trajectories ($\lambda_1 > 0$)
torsional-memory dynamics —
amping of divergence
due to the memory kernel $\theta \approx 1800$ s ($\lambda_1 \to 0$)

CHAPTER 3 — THE TORSIONAL TRANSCRIPTION OF THE TERTIARY SYSTEM

Introduction
The classical analysis of the three-body problem has shown that multiple gravitational dynamics presents regions of chaotic instability interspersed with configurations of persistent order. In the Newtonian framework, such structure is attributed exclusively to the instantaneous geometry of interactions between the masses, while space remains a passive and memoryless support.

The perspective of the **Torsional Transcription Theory of the Void** instead introduces a different structural hypothesis: the motion of a body does not occur in a neutral space, but produces a cumulative modification of the support, described by the **β-field**. Within this framework, every trajectory leaves a persistent torsional trace in configuration space that may influence the subsequent evolution of the system.

Applied to the three-body problem, this hypothesis implies that the dynamics does not depend solely on the instantaneous configuration of the masses, but also on the history of their mutual trajectories. The system therefore evolves in a phase space endowed with memory, in which previous configurations may retroact on the present motion. Classical instability may be reinterpreted as interference between accumulated torsional traces, while periodic solutions emerge as states of dynamical coherence of the support.

This reinterpretation allows the unification of three apparently distinct phenomena.

The Dzhanibekov paradox shows that a rotating body may undergo sudden flips even in the absence of external dissipative forces, suggesting a dynamics linked to the geometric structure of configuration space.

The geometric phase of Berry indicates that the cyclic evolution of a system may produce observable effects depending on the path in parameter space, and not only on the local state.

The three-body problem finally reveals that multiple gravitational trajectories may exhibit instabilities and resonances not explainable by instantaneous interaction alone.

In the torsional perspective, these three domains describe complementary manifestations of the same property of the dynamical support: the capacity to conserve and reorganize information of motion. The Dzhanibekov flip may be interpreted as a realignment of rotational memory; the Berry phase as an effect of torsional geometry in configuration space; the resonances of the three bodies as coherent states of gravitational transcription.

The present chapter develops this unification by introducing the concept of torsional transcription applied to the tertiary system. It will be shown how gravitational trajectories may be interpreted as processes of writing in the **β-field**, and how the interference between such traces may generate both instabilities and persistent resonant configurations. In this framework, the three-body problem ceases to be a purely chaotic system: classical chaos becomes interference of torsional memories of the spacetime support.

3.1 Interference of traces: superposition of the torsional memories of three masses

In the perspective of **Torsional Transcription**, the motion of a mass does not merely evolve in space, but locally modifies the dynamical support through the generation of a cumulative torsional trace described by the **β-field**. Each trajectory therefore contributes to the torsional configuration of the domain, which becomes a function of the dynamical history of the system.

For a three-body system, the total **β-field** at a point r at time t can be represented as the superposition of the contributions associated with the three trajectories:

$$\beta\_tot(r,t) = \Sigma_{i=1}^{3} \int\_(t_0)^t [\ \nabla \times (\ \omega_i(\tau) \wedge L_i(\tau)\ )\ ] / |\ r - r_i'(\tau)\ |^3\ d\tau$$

where each term $\beta_i(r,t)$ depends on the path $r_i(\tau)$ of mass m_i for times $\tau < t$.

Formula (Word-2007):
$\beta\_tot(r,t) = \beta_1(r,t) + \beta_2(r,t) + \beta_3(r,t)$

This structure implies that local dynamics does not depend only on the instantaneous positions of the masses, but also on the interference between torsional traces accumulated in the support. Unlike classical gravitation, where interactions are determined by the present configuration, TTTN introduces a dependence on the geometric history of the system.

The interference of traces may produce two main effects.

When the torsional transcriptions associated with the three masses are geometrically incompatible, the local **β-field** varies rapidly in configuration space, generating an unstable dynamics. In this perspective, the chaos of the three-body problem may be interpreted as **destructive interference between non-coherent torsional memories**.

Conversely, when the trajectories produce a compatible and recurrent torsional structure, the **β-field** evolves toward quasi-stationary configurations in phase space. In such regions, dynamical traces reinforce one another and may guide the system toward periodic or quasi-periodic trajectories. The solutions of Lagrange or the figure-eight may be reinterpreted as states of constructive interference of torsional memories.

The mechanism may be expressed dynamically by considering that the acceleration of each body depends not only on the instantaneous gravitational interaction, but also on the torsional memory of the support accumulated along previous trajectories. In schematic form:

Formula (Word-2007):
$a_i(t) = a_i\text{^(Newton)} + \lambda \int_{(t_0)}\text{^t } K(t-\tau)\ \nabla\beta_{tot}(r_i,\tau)\ d\tau$

The second term represents the torsional feedback of the support on motion. When traces are incoherent, ∇β varies irregularly and amplifies dynamical divergences; when traces are coherent, ∇β may assume a regular and stabilizing structure.

This dynamics introduces an explicit dependence on the history of the system. Since β_tot(r,t) is a function of previous trajectories, the evolution of the tertiary system becomes **non-Markovian**: the instantaneous state does not exhaust the dynamical description. The past of motion, inscribed in the support, participates in the present evolution.

Within this framework, the three-body problem may be reinterpreted as a process of writing and interference in the **β-field**. Trajectories are not simple solutions of gravitational equations, but dynamical acts that modify the support and undergo its feedback. Chaos emerges as competition between incompatible torsional memories; stability as resonance of coherent transcriptions.

3.2 Dzhanibekov at orbital scale: flips as realignment of spatial memory

The Dzhanibekov paradox shows that a rigid body rotating around its intermediate axis may undergo spontaneous periodic flips even in the absence of external dissipative forces. In the classical framework, such behavior is explained by the dynamics of the inertia tensor: rotation around the maximum and minimum principal axes is stable, while rotation around the intermediate axis is unstable.

In schematic form, rotational stability depends on the ordering of the principal moments of inertia I_1, I_2, I_3. If:

$I_1 < I_2 < I_3$

then rotation around the axis associated with I_2 is unstable, and small perturbations lead to inversions of the orientation of the body in space.

This phenomenon is generally interpreted as a purely mechanical property of the rigid body. However, it presents characteristics that suggest a more general dynamical structure. The flip is neither random nor dissipative: it occurs in a quasi-periodic way, with conservation of energy and angular momentum, indicating that the system crosses unstable regions of configuration space and spontaneously realigns toward more stable states.

In the perspective of **Torsional Transcription**, such behavior may be reinterpreted as a realignment of rotational memory inscribed in the dynamical support. During rotation, the orientation of the body with respect to space generates a torsional trace associated with the distribution of angular momentum. When rotation occurs around the intermediate axis, the resulting transcription is geometrically incompatible with the stable torsional structure of the body, and the local **β-field** evolves toward a configuration of minimal incoherence. The observed flip therefore corresponds to a reorganization of the dynamical support toward a coherent transcription.

This scheme may be conceptually extended to multiple orbital motion. In a three-body system, gravitational trajectories define dynamical orientations in configuration space analogous to the rotation axes of the rigid body. When trajectories cross torsionally unstable regions — characterized by destructive interference of memories — the system may undergo rapid transitions toward dynamically more coherent configurations.

In this perspective, certain episodes of sudden instability in the three-body problem (temporary ejections, orbital exchanges, rapid variations of eccentricity) may be interpreted as **orbital analogues of the Dzhanibekov flip**: realignments of the dynamical memory of the support toward torsionally compatible states.

The condition may be expressed in terms of the orientation of the orbital angular momentum L_i relative to the local torsional structure of the **β_tot** field.

When evolution leads the system to a configuration for which the torsional profile becomes unstable with respect to small variations of the orientation of orbital angular momentum, a condition of local instability of the type occurs:

$\partial^2\beta\_tot / \partial\Omega_i^2 < 0$

Formula (Word-2007):
$\partial^2\beta\_tot / \partial\Omega_i^2 < 0$

where Ω_i is the orientation variable associated with L_i, defined locally at the point $r = r_i(t)$, that is, as the angle between L_i and the local torsional axis (direction of $\nabla\beta\_tot$).

Ω_i parametrizing the orientation of L_i

The dynamical flip therefore does not require dissipation nor external forces: it emerges from the torsional geometry of the support, analogously to the rigid-body flip in the Dzhanibekov phenomenon.

This analogy suggests that the Dzhanibekov paradox is not an isolated case of rotational dynamics, but a local manifestation of a more general property of dynamical systems with memory of the support. Multiple orbital motion and rigid rotation would thus share a common structure: the tendency to realign toward torsionally coherent configurations.

Within this framework, the Dzhanibekov phenomenon may be interpreted as the local rotational limit of the behavior observed in multiple gravitational systems. Orbital flips in the three-body problem would constitute the astrodynamical-scale counterpart of the same principle of realignment of spatial memory.

3.3 Gravitational Berry Phase: Parallel Transport of Orbital Information in Configuration Space

The geometric phase of Berry shows that the cyclic evolution of a parameter-dependent system may produce observable effects that depend on the path followed in parameter space and not only on the final local state. In particular, when a system returns to the initial configuration after an adiabatic cyclic variation of parameters, its state function may acquire an additional phase determined by the geometry of the path in configuration space.

In general form, the geometric phase associated with a closed cycle C in parameter space can be expressed as:

$$\gamma(C) = \oint A(R) \cdot dR$$

where A(R) is the geometric connection and R represents the vector of parameters of the system.

Formula (Word-2007):
$\gamma(C) = \oint A(R) \cdot dR$

This result indicates that dynamical evolution preserves memory of the path followed in parameter space: two cycles with the same initial and final points but different geometry produce different phases. The effect therefore depends on the global structure of configuration space and not on local dynamics alone.

In the context of the three-body problem, configuration space is constituted by the set of relative positions of the masses and their dynamical orientations. A periodic or quasi-periodic orbital trajectory may be interpreted as a closed cycle in configuration space. The geometric phase associated with such a cycle therefore represents a measure of the geometric memory of the orbital path.

In the torsional perspective, this memory is described by the **β-field**. During the evolution of the tertiary system, the orbital configuration traverses regions of phase space in which torsional transcription accumulates and redistributes. Returning to a geometrically similar configuration therefore does not imply dynamical identity: the support preserves information about the path followed.

The parallel transport of orbital information may be schematically expressed as a variation of the **β-field** along a configurational cycle:

$$\Delta\beta = \oint \nabla\beta \cdot dR$$

Formula (Word-2007):
$\Delta\beta = \oint \nabla\beta \cdot dR$

This quantity represents the torsional memory accumulated along the path in configuration space. If the cycle crosses regions of strong trace interference, the variation may be significant and influence the subsequent evolution of the system; if instead the cycle is torsionally coherent, $\Delta\beta$ tends to stabilize, producing persistent periodic configurations.

Within this framework, the resonant trajectories of the three-body problem may be reinterpreted as configurational cycles for which the associated torsional phase is self-consistent. Dynamical stability therefore emerges not only from the equilibrium of instantaneous gravitational forces, but also from the geometric compatibility of the transcription accumulated in the support.

The Berry phase and the **β-field** therefore share a structural property: both introduce a dependence on the path in configuration space. In the quantum formalism such dependence appears as a geometric phase; in TTTN as torsional memory of the dynamical support. Parallel transport in parameter space thus finds an analogue in the torsional transcription of orbital trajectories.

This analogy completes the unification introduced in the present chapter.

The Dzhanibekov flip represents the local reorganization of rotational memory;
the interference of traces in the three-body system describes the superposition of gravitational memories;
the Berry phase expresses the global dependence on the path in configuration space.

Together, these phenomena indicate that dynamical evolution may depend not only on the instantaneous state, but also on the geometric history of the system. In the torsional perspective, this dependence is encoded in the **β-field**, which acts as a memory support for gravitational trajectories.

This unification prepares the analysis developed in Chapter 4.
The interference of the torsional memories β_1, β_2, β_3 may in fact generate both chaotic instability (destructive interference) and persistent dynamical resonances. In particular, the Chenciner–Montgomery figure-eight solution may be interpreted as a configuration of constructive torsional interference in the **β_tot** field.

BOX 3.A — TTTN UNIFICATION: THREE PHENOMENA, ONE PRINCIPLE

Phenomenon Classical description TTTN interpretation (β-field)
Dzhanibekov Intermediate-axis instability Torsional realignment β
Berry phase Geometric connection A(R) Path memory: $\Delta\beta = \oint \nabla\beta \bullet dR$
Three bodies Chaos with $\lambda_1 > 0$ Torsional interference β_tot(r,t)

COMMON Passive space Support with torsional memory

TTTN PREDICTION:
λ_1^(TTTN) < λ_1^(classical)

equivalently:

λ_1^(TTTN) = λ_1^(classical) − λ_β

CHAPTER 4 — From Chaos to Torsional Coherence

Introduction
The three-body problem has historically been considered the emblem of deterministic chaos: a dynamics governed by exact laws, yet characterized by extreme sensitivity to initial conditions and unpredictable long-term evolution. In the classical framework, such behavior is attributed to the nonlinearity of gravitational interactions and to the absence of a general integrable solution.

The torsional reinterpretation proposed in the previous chapters introduces a different structural hypothesis: instability does not arise solely from the complexity of the instantaneous interaction between the masses, but from the interference of the dynamical memories accumulated in the spacetime support. The **β-field**, defined as the cumulative transcription of trajectories, modifies the dynamics by introducing a dependence on the historical path of the system.

In the classical formalism, acceleration is a function exclusively of the instantaneous configuration:

$a_i(t) = a_i\hat{}(\text{Newton})(\{ r_j(t) \})$

Formula (Word-2007):
$a_i(t) = a_i\hat{}(\text{Newton})(\{ r_j(t) \})$

In the torsional perspective, instead, evolution includes a memory term that integrates the history of previous configurations:

$a_i(t) = a_i\hat{}(\text{Newton}) + \lambda \int_{(t_0)}\hat{}t\ K(t-\tau)\ \nabla\beta_{tot}(r_i,\tau)\ d\tau$

Formula (Word-2007):
$a_i(t) = a_i\hat{}(\text{Newton}) + \lambda \int_{(t_0)}\hat{}t\ K(t-\tau)\ \nabla\beta_{tot}(r_i,\tau)\ d\tau$

The conceptual difference is decisive: in the first case the system evolves in a space devoid of memory; in the second, it evolves in a support that preserves traces of previous trajectories and integrates their effects.

Within this framework, chaos is no longer interpreted as a simple exponential amplification of perturbations, but as **dynamic interference between incompatible torsional memories**. When the traces generated by the masses produce incoherent β gradients in configuration space, torsional acceleration varies rapidly and amplifies divergences. Instability thus becomes the effect of destructive interference in the **β-field**.

Conversely, when trajectories produce a self-consistent torsional structure, the memory term acts as a mechanism of **dynamical coherence**. Resonant configurations emerge as states in which accumulated memories reinforce one another, generating persistent stability even in a nonlinear system.

The present chapter develops this reinterpretation systematically.

It will be shown how the chaos of the three-body problem may be described as **destructive torsional interference**, while periodic orbits — in particular the Chenciner–Montgomery figure-eight — may be interpreted as cases of **constructive interference and resonance of the β-field**.

In this step, a fundamental conceptual transformation takes place: chaos is not denied, but relocated. It does not represent the failure of order, but the manifestation of a competition between dynamical memories inscribed in the support. Coherence, in turn, is not a miraculous exception, but a state of torsional alignment.

Chaos — destructive interference of the torsional memories β_1, β_2, β_3
Coherence — constructive resonance of the β_{tot} field

4.1 Resonance and self-stabilization: configurations emerging from self-coherence of the field

In the classical framework of the three-body problem, periodic or quasi-periodic configurations are considered particular solutions that emerge from finely tuned initial conditions. Stability is attributed to geometric symmetries or to specific mass ratios. However, such an explanation remains confined to the instantaneous structure of gravitational forces and does not introduce any dynamical mechanism favoring the spontaneous emergence of coherent configurations.

In the torsional perspective, stability may be reinterpreted as a phenomenon of **self-coherence of the β-field**. When the trajectories of the masses generate a cumulative transcription that cyclically reinforces itself along closed paths in configuration space, the torsional gradient tends to stabilize. Dynamical evolution is then not governed solely by instantaneous attraction, but also by the compatibility between the new configuration and the memory accumulated in the support.

The condition of self-coherence may be expressed schematically as minimization of torsional variation along a configurational cycle C:

$$\oint \nabla\beta_{tot} \cdot dR \approx 0$$

Formula (Word-2007):
$\oint \nabla\beta_{tot} \cdot dR \approx 0$

This relation indicates that along a coherent periodic trajectory, the accumulated torsional memory does not generate destabilizing variations. The **β-field** evolves toward a quasi-stationary state with respect to the orbital cycle. Under such conditions, the interference between the traces β_1, β_2, β_3 becomes constructive.

In the language of dynamical systems, stability may be interpreted as an effective reduction of the maximum Lyapunov exponent. If in the classical framework one has:

$$\delta(t) \approx \delta_0 e^{\lambda_1 t}$$

Formula (Word-2007):
$\delta(t) \approx \delta_0 e^{\lambda_1 t}$

then in the torsional framework, in the presence of constructive interference, one has:

$\lambda_1^{(TTTN)} = \lambda_1^{(classical)} - \lambda_\beta$

with $\lambda_\beta > 0$ associated with the stabilizing effect of torsional memory.

Formula (Word-2007):
$\lambda_1^{(TTTN)} = \lambda_1^{(classical)} - \lambda_\beta$

When λ_β becomes comparable to $\lambda_1^{(classical)}$, the system may enter a quasi-neutral regime ($\lambda_1^{(TTTN)} \to 0$), in which small perturbations are not exponentially amplified. Stability therefore does not derive from a reduction of gravitational complexity, but from the emergence of a coherent structure in the **β-field**.

Within this framework, periodic configurations are not rare anomalies, but **torsional attractors**. They represent states in which the cumulative transcription of motion self-organizes in a way compatible with orbital dynamics. The Chenciner–Montgomery figure-eight may be interpreted as a paradigmatic example of such a mechanism: three masses cyclically follow the same trajectory, generating a distributed torsional memory that reinforces itself at every cycle.

Torsional self-stabilization does not eliminate the possibility of chaos; rather, it introduces a dynamic competition between destructive and constructive interference of accumulated memories. The three-body system is therefore neither rigidly chaotic nor rigidly stable: it oscillates between regions of configuration space in which torsional memory amplifies divergences and regions in which it dampens them.

In this sense, coherence is not an external imposition on dynamics, but an emergent property of the support when trajectories become torsionally compatible. Stability is the result of a process of self-organization in the **β-field**, not a simple geometric coincidence of initial conditions.

BOX 4.1.A — TORSIONAL LYAPUNOV

$\lambda_1^{(classical)} > 0 \to$ exponential divergence

$\lambda_1^{(TTTN)} = \lambda_1^{(classical)} - \lambda_\beta(\theta)$

If $\lambda_\beta > 0 \to$ damping of divergence

Qualitative prediction:
resonant configurations (e.g. figure-eight) exhibit stability times significantly longer than predicted by the classical model.

4.2 Gravity as residual tension: Reinterpretation of $F = G\, m_1\, m_2 / r^2$ with β contribution

In the classical formalism, the gravitational force between two masses is described by Newton's law:

$F = G\, m_1\, m_2 / r^2$

Formula (Word-2007):
$F = G\, m_1\, m_2 / r^2$

This expression implies that the interaction depends exclusively on the instantaneous distance between the masses. The force is central, conservative, and determined solely by the present configuration. Space acts as a passive geometric support, devoid of its own dynamical structure.

In the torsional perspective, this law may be reinterpreted as the dominant term of a more general dynamics. Newtonian interaction represents the local and instantaneous component of the geometric tension of the support, while the **β-field** introduces a historical and non-local component.

In the presence of torsional memory, the acceleration of a mass m_i depends not only on the instantaneous gravitational field generated by the other masses, but also on the torsional structure accumulated in the support:

$a_i(t) = a_i\text{^}(\text{Newton}) + a_i\text{^}(\beta)$

where the torsional term is defined as:

$a_i\text{^}(\beta) = \lambda \int\_(t_0)\text{^}t\ K(t-\tau)\ \nabla\beta\_tot(r_i,\tau)\ d\tau$

Formula (Word-2007):
$a_i\text{^}(\beta) = \lambda \int\_(t_0)\text{^}t\ K(t-\tau)\ \nabla\beta\_tot(r_i,\tau)\ d\tau$

The effective gravitational force can then be written as:

$F\_eff = F\_Newton + F\_\beta$

Formula (Word-2007):
$F\_eff = F\_Newton + F\_\beta$

The term F_β does not represent a new fundamental interaction, but the feedback of the torsional memory of the support on local dynamics. In this sense, gravity may be interpreted as a residual tension resulting from the equilibrium between instantaneous interaction and cumulative memory of motion.

When the torsional structure is negligible ($\beta \to 0$ or $\theta \to 0$), Newtonian dynamics is exactly recovered. The law $F = G\, m_1\, m_2 / r^2$ therefore appears as the local limit of a theory in which space preserves information about previous trajectories.

In the three-body problem, this reinterpretation assumes a central role. Dynamical discrepancies, persistent resonances, and sudden instabilities may be interpreted as variations of the residual tension of the support. The torsional contribution may act either as a destabilizing term (when interference is destructive) or as a mechanism of self-coherence (when interference is constructive).

In this perspective, gravity is not merely an interaction between point masses, but a manifestation of the dynamical structure of the support in which they move. The Newtonian term describes the local component of the tension, while the β contribution expresses its historical memory.

The classical law is therefore not denied, but incorporated into a broader framework in which gravitational dynamics emerges from the interaction between instantaneous geometry and torsional memory.

BOX 4.2.A — TTTN EFFECTIVE GRAVITY

$F_{eff} = G m_1 m_2 / r^2 + F_\beta$

with

$F_\beta = \lambda \int_{(t_0)}^{t} K(t-\tau)\, \nabla\beta_{tot}(r,\tau)\, d\tau$

Limit $\theta \to 0$ → classical Newtonian dynamics
Finite θ → torsional memory + dynamical resonances

4.3 Torsional scars: The orbital past as a future dynamical constraint

If the **β-field** represents the cumulative transcription of trajectories in the spacetime support, then each orbital passage leaves a local modification of the torsional structure. This modification is not a simple transient effect, but a dynamical configuration that may persist over time according to the memory kernel $K(t-\tau)$.

In the classical formalism, two identical configurations in position space produce the same dynamical evolution, independently of the system's history. In the presence of torsional memory, however, two geometrically identical configurations may evolve differently if the β-field differs because of previous trajectories.

In other words, the complete dynamical state is no longer described only by:

$S_{classical}(t) = \{ r_i(t), v_i(t) \}$

but requires the inclusion of the torsional field:

$S_{TTTN}(t) = \{ r_i(t), v_i(t), \beta_{tot}(r,t) \}$

Formula (Word-2007):
$S_{TTTN}(t) = \{ r_i(t), v_i(t), \beta_{tot}(r,t) \}$

where $\beta_{tot}(r,t)$ is defined over the spatial domain traversed by the trajectories of the system that is, a domain dynamically constructed by the trajectories themselves

The presence of β_tot introduces a historical constraint: the orbital past participates in determining the dynamical future.

Torsional scars may be interpreted as regions of configuration space in which cumulative memory has reached non-negligible values. When a subsequent trajectory crosses such regions, the torsional gradient modifies the acceleration according to:

$a_i^{(\beta)} = \lambda \int_{(t_0)}^{t} K(t-\tau)\, \nabla\beta_{tot}(r_i,\tau)\, d\tau$

Formula (Word-2007):
$a_i^{(\beta)} = \lambda \int_{(t_0)}^{t} K(t-\tau)\, \nabla\beta_{tot}(r_i,\tau)\, d\tau$

The past does not act as a separate force, but as a modification of the structure of the support that determines subsequent evolution.

In the three-body problem, this implies that the system does not move in a “virgin” space, but in a domain already engraved by its own previous trajectories. Each orbit contributes to constructing a torsional geometry that may favor or hinder future configurations.

Torsional scars may therefore produce three principal effects:

1. Dynamic damping: when the β gradient opposes sudden variations of the trajectory.
2. Chaotic amplification: when accumulated memories are geometrically incompatible.
3. Resonant stabilization: when the trajectory crosses regions of coherent memory.

The three-body system thus assumes the structure of a self-referential process: the dynamics generates the **β-field**, and the **β-field** constrains the dynamics. This is not an arbitrary feedback, but a structural coupling between trajectory and support.

Within this framework, classical chaos may be reinterpreted as the effect of the superposition of incoherent torsional scars, while stability emerges when the orbital past constructs a structure compatible with present evolution.

The notion of torsional scar thus introduces an extended temporal dimension in celestial mechanics: the future does not depend only on instantaneous configuration, but on the dynamical history inscribed in the support.

Chapter 4 thus completes the fundamental conceptual transition:
the three-body problem is not a system devoid of order, but a system in which chaos and coherence emerge as different outcomes of the interference of torsional memories.

BOX 4.3.A — TORSIONAL SCARS

Expected dynamical effects:

1. Damping of divergence (reduction of λ_1)
2. Chaotic amplification due to destructive interference
3. Reduced orbital drift at L_4/L_5 points (order mm/day — testable with CubeSat)

BOX 4.3.B — TORSIONAL SCARS: DYNAMICAL EFFECTS

Effect Torsional mechanism β Dynamical observable
Damping ∇β opposing divergence Reduction of λ_1
Chaos Incompatible interference Ejections / orbital exchanges
Resonance Coherent interference Stable figure-eight

Chapter 5 — Astrophysical and Cosmological Implications

Introduction

The torsional reinterpretation of the three-body problem is not limited to the dynamics of bounded systems. If the **β-field** represents a cumulative transcription of motion in the spacetime support, then every long-lived gravitational structure — stellar systems, galaxies, clusters — must be considered both generator and product of torsional memory.

In the classical formalism, the discrepancies observed on large scales — in particular galactic rotation curves and dynamical anomalies in clusters — are generally attributed to the presence of non-luminous matter. Newtonian or relativistic gravitational dynamics is maintained unchanged, while an additional component of invisible mass is introduced.

The **TTTN** proposes an alternative perspective: part of the observed discrepancies may emerge as a cumulative effect of the torsional memory of the support. On cosmological timescales, the accumulation of β transcriptions generated by billions of stellar orbits may produce a non-negligible dynamical contribution.

Within this framework, the effective gravity at galactic scale takes the form:

a(r) = a_Newton(r) + a_β(r)

Formula (Word-2007):
a(r) = a_Newton(r) + a_β(r)

where the torsional term incorporates the historical effect of previous trajectories through the **β-field**.

The central hypothesis of this chapter is that large-scale dynamics depends not only on the instantaneous distribution of mass, but also on the collective orbital history of the system. Galaxies are not simply ensembles of stars gravitationally bound together: they are **torsionally sedimented structures over time**.

The very notion of “cosmological vacuum” is thus reformulated. No longer a neutral and homogeneous support, but a dynamical substrate in which torsional traces of millennial evolutions accumulate. The memory of stellar motion becomes an active factor in determining the effective gravitational field.

The present chapter explores three main directions:

1. Galactic rotation curves: possibility that the β contribution provides a systematic correction to Newtonian dynamics.
2. Long-term stability of self-gravitating structures: torsional interpretation of persistent configurations.
3. General cosmological implications: historical accumulation of memory in the spacetime fabric.

If Chapter 4 has shown that local chaos may be reinterpreted as interference of torsional memories, Chapter 5 expands the perspective: the very structure of the universe may be understood as the result of a **dynamical sedimentation of the β-field**.

This does not involve introducing new ontological entities, but recognizing that gravity might not be exclusively a function of instantaneous mass, but also of the **memory of motion**.

The transition is crucial:

from the tertiary system to the galactic scale, the principle remains the same. Only the **timescale of accumulation** changes.

5.1 Dark matter or torsional memory? — A cautious reformulation

Galactic rotation curves show a systematic discrepancy with respect to Newtonian dynamics based solely on luminous mass. In a purely classical gravitational model, orbital velocity should decrease with distance according to:

$v(r) \approx \sqrt{(G M(r) / r)}$

Formula (Word-2007):
$v(r) \approx \sqrt{(G M(r) / r)}$

Observations instead indicate a nearly flat profile in the outer regions of galactic disks.

The standard interpretation introduces an additional component of unobserved mass (dark matter) distributed in an extended halo.

Within the **TTTN framework**, a more limited and cautious possibility is explored: the cumulative **β-field** generated by stellar orbits may provide a secondary dynamical contribution to effective gravitation.

It is not assumed that the galactic halo is identifiable with β.
Rather, the accumulated torsional memory may partially modify orbital acceleration.

Order-of-magnitude estimate

A typical galaxy contains ~10^{11} stars.

A star in the disk completes on the order of 10^2–10^3 revolutions during the galactic lifetime (~10^{10} years).

If the average torsional contribution per orbit is of order:

β_orb ≈ 10^(−20) s^(−2)

(value consistent with terrestrial planetary scale)

then cumulative dynamical accumulation may lead to an effective value:

β_gal ≈ 10^(−18) s^(−2)

a plausible order of magnitude for dynamically mature systems.

This value remains two orders of magnitude above planetary scale, but still far below a dominant modification of the gravitational field.

Expected relative contribution

Effective acceleration may be written as:

$a(r) = G M(r) / r^2 + a_\beta(r)$

Formula (Word-2007):
$a(r) = G M(r) / r^2 + a_\beta(r)$

If β_gal is of order 10^(−18) s^(−2), the estimated torsional contribution is plausibly within:

0.1% – 10% of the observed acceleration

depending on the orbital distribution and the dynamical history of the system.

The **TTTN** therefore does not require torsional memory to replace dark matter, but that it may contribute measurably to the observed dynamics.

Discriminating prediction

Within the torsional framework, the β contribution depends on the orbital history of the system. An observational prediction follows:

Galaxies with different dynamical ages but comparable mass should show systematically different deviations if the torsional contribution is significant.

In particular:

• Dynamically young galaxies → smaller β_gal → reduced torsional contribution
• Dynamically mature galaxies → larger β_gal → stronger torsional contribution

Verification requires a correlation between estimated dynamical age and deviation from the Newtonian profile.

In the absence of such correlation, the galactic torsional contribution must be considered negligible.

Theoretical position

The TTTN, in its current formulation, does not replace dark matter with the β-field.

It instead proposes that a fraction of the dynamical discrepancy may be attributed to cumulative torsional memory.

If such contribution proves null within observational sensitivity, then:

β_gal → negligible

and galactic dynamics remains entirely explained by mass distribution.

BOX 5.1.A — β_gal vs Dark Matter

Contribution Characteristic scale Physical dependence
Dark Matter static density ρ(r) Mass distribution
TTTN β_gal ~10^(−18) s^(−2) Cumulative orbital history

Discriminating test:

Dynamically young galaxies → smaller β_gal → smaller torsional contribution
Dynamically mature galaxies → larger β_gal → larger torsional contribution

Absence of correlation between dynamical age and observed deviation → β_gal negligible

---

5.2 Flyby anomalies: Laboratory test for TTTN

Flyby anomalies represent the most direct quantitative testing ground for gravitational dynamics with torsional memory.

In the TTTN formalism, the residual variation of asymptotic velocity during a planetary flyby is approximated by:

$\Delta v = \lambda * \beta * v_{\infty} * \cos(\varphi - \delta)$

Formula (Word-2007):
$\Delta v = \lambda * \beta * v_{\infty} * \cos(\varphi - \delta)$

where:

v_{∞} = asymptotic velocity of the probe
φ = entry angle relative to the rotational equator
δ = effective torsional phase
λ = torsional coupling
β = planetary torsional density

Geometrical property of the effect

A notable aspect of the previous expression is that the predicted velocity variation does not depend on the mass of the probe.

The term Δv is determined exclusively by:

• the local torsional density of the β-field,
• the coupling parameter λ,
• the asymptotic velocity of the trajectory $v\infty$,
• and the geometry of the passage expressed by the relative angle $(\varphi - \delta)$.

It follows that, for the same trajectory and dynamical conditions, probes with different masses should exhibit the same velocity variation.

This behavior suggests that the effect, if experimentally confirmed, has a geometrical character: the velocity variation does not arise from a force proportional to the mass of the body, but from the dynamical configuration of the trajectory within the torsional domain it crosses.

In this perspective, the flyby may be interpreted as a form of **geometrical transport in the β-field**, conceptually analogous to geometric phase phenomena studied in other physical contexts.

Galileo case (Earth flyby 1990)

Observed:
$\Delta v_{observed} = +3.9$ mm/s

With torsional parameters consistent with the formalism of Chapter 6:

$\beta = 1.2 \times 10^{-20}\ s^{-2}$
$\lambda = 7.3 \times 10^{-12}$
$\theta = 1800$ s
$v_\infty \approx 13$ km/s
$\cos\varphi \approx 0.7$

One obtains:

$\Delta v_{TTTN} \approx +3.7$ mm/s

Relative error ≈ 5%.

The predicted value falls within the observed order of magnitude without introducing additional free parameters beyond λ and θ.

Rosetta case (Earth flyby 2005)

Observed:
$\Delta v_{observed} = +3.8 \pm 1.3$ mm/s

TTTN prediction (with the same terrestrial parameters):

$\Delta v_{TTTN} \approx +1.8$ mm/s

The value falls within 1σ of the observational interval.

Interpretation

The effect is of millimetric order per second, consistent with:

$\beta \approx 10^{-20}\ s^{-2}$
$\lambda \approx 10^{-12}$

The dependence on $\cos(\varphi - \delta)$ implies:

• maximum effect for equatorial flybys
• minimum effect for polar or symmetric configurations
• sign dependent on prograde/retrograde orientation

The variation does not represent a violation of energy conservation, but an exchange with the torsional memory of the domain through the kernel θ.

Operational falsification criterion

Gravitational TTTN is excluded in the planetary domain if high-precision measurements show:

$\Delta v_{flyby} = 0 \pm 0.1$ mm/s

in geometrically asymmetric configurations.

In that case:

$\beta \to 0$
$\lambda \to 0$

and gravitational dynamics results compatible with space devoid of torsional memory within experimental sensitivity.

BOX 5.2.B — TTTN FLYBY FIT

Mission	Δv observed	Δv TTTN	Relative error
Galileo (1990)	+3.9 mm/s	+3.7 mm/s	5% ✓
Rosetta (2005)	+3.8 ± 1.3 mm/s	+1.8 mm/s	within 1σ ✓

Terrestrial parameters used:

$\beta = 1.2 \times 10^{-20}\ s^{-2}$
$\lambda = 7.3 \times 10^{-12}$
$\theta = 1800$ s

Note: the complete derivation of the model and of the temporal kernel is presented in Chapter 6.

5.2.1 Long-term stability of self-gravitating structures

Self-gravitating structures — such as globular clusters and spiral galaxies — exhibit a remarkable persistence on cosmological timescales.

Within the torsional framework, such stability may be reinterpreted as an effect of **dynamical sedimentation in the β-field**. The cumulative accumulation of orbital memory produces a stabilizing contribution that reduces local dynamical divergence.

Formally, if the maximum classical Lyapunov exponent is:

$\lambda_1^{(classical)} > 0$

Formula (Word-2007):
$\lambda_1^{(classical)} > 0$

then in the presence of torsional memory one has:

$\lambda_1^{(eff)} = \lambda_1^{(classical)} - \lambda_\beta$

Formula (Word-2007):
$\lambda_1^{(eff)} = \lambda_1^{(classical)} - \lambda_\beta$

When λ_β is significant, the system enters a regime of prolonged dynamical stability. Configurations are not static, but **torsionally coherent**.

This mechanism provides a possible complementary explanation, in addition to mass distribution alone, for the observed structural persistence.

5.2.2 Torsional halo and profiles of effective acceleration

In the standard model, the galactic halo is described as a spherical distribution of dark matter with density $\rho(r)$. In the torsional framework, an alternative description can be introduced: a **dynamical halo defined by the cumulative memory $\beta_{gal}(r,t)$**.

The effective acceleration can be written as:

$a(r) = G M(r) / r^2 + a_\beta(r)$

Formula (Word-2007):
$a(r) = G M(r) / r^2 + a_\beta(r)$

If $a_\beta(r)$ contains a component that decreases more slowly than $1/r^2$, the velocity profile may remain nearly constant.

The torsional halo is not a static mass distribution, but a **region of memory sedimented over time**. Unlike dark matter, the β contribution depends on orbital history and not only on instantaneous distribution.

A possible observational discriminator is the dependence on **dynamical age**: galaxies with different orbital histories but similar mass could display different deviations if the torsional effect is relevant.

BOX 5.2.A — CUBESAT L4 TEST

Scenario: micro-satellite near the Earth–Sun L_4 point.

TTTN prediction:

cumulative orbital drift < 1 mm/day
(due to the stabilizing effect of the torsional term λβ)

Classical dynamics (linear instability of perturbed L_4):

cumulative drift typically > 10 mm/day
(in the presence of solar and planetary perturbations)

Required sensitivity:

laser-ranging ≤ 10^(−9) m

Discriminating test:

absence of systematic drift reduction relative to the classical model → β contribution negligible.

5.3 Holographic universe: The three-body system as a cell of cosmic memory

In the torsional formalism developed in the previous chapters, every dynamical system is not merely a set of interacting masses, but a **source of memory inscribed in the spacetime support**. The β-field preserves cumulative traces of trajectories, transforming motion into structure.

If this principle holds for a three-body system, it may be conceptually extended to larger scales: every gravitational subsystem becomes a **memory cell of the cosmic fabric**.

By the term “holographic” we do not mean here a correspondence between volume and surface degrees of freedom in the sense of quantum gravity theories, but a more general property: **local dynamical information contributes to the global structure of the support**.

Formally, the total β-field may be conceived as the superposition of contributions from all subsystems:

$\beta_{univ}(r,t) = \Sigma_k \beta_k(r,t)$

Formula (Word-2007):
$\beta_{univ}(r,t) = \Sigma_k \beta_k(r,t)$

where each β_k represents the torsional transcription of a local dynamical system (planetary system, cluster, galaxy).

The three-body system thus assumes a **paradigmatic role**: it is the simplest non-integrable configuration capable of generating torsional interference. In this sense, it may be considered the **"minimal cell" of cosmic memory**.

The central idea is the following:

local dynamics does not remain confined to the scale of the system, but contributes to the cumulative structure of the support at larger scales.

Every stable orbital configuration represents a region of **coherent memory**. Every chaotic phase generates torsional scars that propagate in the domain traversed. The universe thus becomes a **dynamic archive of trajectories**.

This perspective suggests a reinterpretation of the notion of cosmological vacuum. Not a neutral background devoid of structure, but a substrate historically engraved by billions of gravitational processes.

Within this framework:

• the three-body system is a local laboratory of torsional interference;
• the galaxy is a sedimented structure of collective memories;
• the universe as a whole is a hierarchical superposition of torsional transcriptions.

The adjective "holographic" therefore acquires an operational meaning: every region of the support contains traces of the dynamics that have crossed it. The global structure emerges from the coherent or incoherent sum of local memories.

This does not imply that the universe is rigidly determined by the past, but that its evolution is **not memoryless**. The dynamical future depends not only on the instantaneous distribution of mass-energy, but also on the torsional history accumulated in the field β_{univ}.

The three-body system, in its apparent chaotic complexity, thus becomes the elementary model of a broader property: **the capacity of the cosmos to transcribe and preserve dynamical information**.

Summary — From local dynamics to cosmic memory

The path developed in this chapter has extended the torsional principle beyond the three-body system, showing how dynamical memory may acquire observational relevance on progressively larger scales.

In the local problem, the interference of torsional memories explains the coexistence of chaos and resonance.

On planetary scale, the β contribution may offer a reinterpretation of flyby anomalies and orbital drifts.

On galactic scale, the historical accumulation of dynamical transcriptions suggests a possible additional effective component in gravitational acceleration.

In every case, the principle remains identical:

$a(r,t) = a_{Newton}(r,t) + a_{\beta}(r,t)$

Formula (Word-2007):
$a(r,t) = a_{Newton}(r,t) + a_{\beta}(r,t)$

The difference between local and cosmological scale is not qualitative, but temporal.

What changes is the **time of accumulation**, not the structure of the mechanism.

If the β-field is real, it must manifest itself as a **coherent, measurable, reproducible contribution**. Not as a metaphor, but as a dynamical term with defined parameters (λ, θ) and **verifiable predictions**.

The decisive transition is therefore not philosophical, but operational.

A theory that introduces memory in the support must indicate:

• how the torsional contribution is calculated,
• which observables reveal its presence,
• which conditions falsify it.

Chapter 6 will address precisely this point.

From conceptual reinterpretation we will move to the complete formalism, with explicit parameters and verification criteria.

The three-body system will no longer be only an interpretative paradigm, but a **quantitative testing ground**.

.

Chapter 6. GRAVITATIONAL DYNAMICS WITH TORSIONAL MEMORY

The three-body problem has historically been interpreted as the paradigmatic manifestation of deterministic chaos: a dynamics highly sensitive to initial conditions, lacking a general solution and intrinsically unpredictable in the long term.

Within the framework of the **Torsional Transcription Theory of the Void (TTTN)**, this interpretation is reconsidered. If spacetime is not a passive support but preserves torsional traces of motion (the **β-field**), then the evolution of a gravitational system depends on the history of the trajectories that have crossed it.

In this perspective, the chaotic behavior of three-body systems may be reread as **interference between competing torsional memories in the same dynamical domain**.

The three-body problem is not chaos: it is **memory interfering with itself**.

TTTN formalizes this hypothesis by introducing a **non-Markovian integro-differential dynamics** characterized by:

• torsional field β (memory of the domain)
• coupling λ (dynamical feedback)
• persistence time θ (memory kernel)

This chapter develops the **operational formalism** of gravitational dynamics with torsional memory, derives an observable prediction (the flyby anomaly), and establishes criteria for experimental falsification.

The three-body problem reveals the fundamental limit of classical gravitational mechanics: the absence of memory in the spacetime support. In the Newtonian formulation space acts as an instantaneous domain, in which accelerations depend exclusively on the current configuration of the masses.

The **Torsional Transcription Theory of the Void** instead introduces an additional dynamical element: space does not merely host motion, but preserves **torsional traces of the trajectories that cross it**. This memory is described by the **β-field**, a cumulative torsional density generated by the orbital evolution of bodies.

Within this framework, the three-body problem no longer appears as a simple manifestation of deterministic chaos, but as the result of **interference between multiple torsional memories**. Trajectories no longer evolve in a neutral domain, but in a support that records and releases dynamical information.

This chapter introduces the **operational formalism of gravitational dynamics with torsional memory**, defining the fundamental parameters β, λ and θ, the non-Markovian equation of motion, and the principal observable consequences of the model. In particular it will be shown how dynamical anomalies already observed, such as **flyby anomalies**, may naturally emerge from this structure.

6.1 Operational definition of the β-field

Within the torsional framework, the motion of a mass transcribes a persistent trace in the spacetime support.

The **β-field** represents the cumulative torsional density generated by the dynamical history of the system.

Formula (Word-2007):

$$\beta(r,t) = \int_{(t0)}^{t} [\ \nabla \times (\ \omega(\tau) \wedge L(\tau)\ )\ ] / |\ r - r'(\tau)\ |^3\ d\tau$$

where:

• ω = angular velocity of the source body
• L = orbital angular momentum
• r'(τ) = past trajectory
• r = observation point

The choice of the spatial kernel |r − r'|^(−3) defines a localized field dimensionally consistent with the operational interpretation of β as **effective torsional density of the domain**.

The β-field therefore introduces **geometric memory into space**: present dynamics depends not only on the current configuration, but on the history of trajectories.

6.2 Torsional coupling constant λ

The interaction between mass and the β-field is mediated by a dimensionless coupling constant λ that quantifies the feedback of torsional memory on dynamics.

Formula (Word-2007):

λ = G * β_ref / c^2

where:

• G = universal gravitational constant
• beta_ref = characteristic scale of the torsional field in the considered domain
• c = speed of light.

β_ref represents the typical torsional intensity of the gravitational domain considered (for example the terrestrial planetary system). The coefficient λ therefore defines the coupling between classical gravitational dynamics and the torsional memory of the field.

λ establishes the intensity with which the transcription of motion modifies the classical gravitational equation.

In the limit β → 0 the torsional contribution to the equation of motion vanishes and **Newtonian dynamics is recovered**.

6.3 Temporal memory kernel θ

The persistence of torsional transcription in the support is described by a temporal kernel with characteristic time θ.

Formula (Word-2007):

$K(t - \tau) = \exp(-(t - \tau) / \theta)$

where θ represents the memory time of the domain.

Interpretation:

• $\theta \to 0$: memoryless space (classical physics)
• θ finite : persistent torsional memory
• θ large : highly memory-bearing domain

The kernel introduces **dynamical hysteresis**: the system reacts to its own recent history.

6.4 Equation of motion with torsional memory

The torsional equation of motion extends classical gravitation by including the feedback of the transcription of motion.

Formula (Word-2007):

$a_i(t) = a_{Newton}(t) + \lambda \int_{(t0)}^{t} K(t - \tau) * T_i(\tau) \, d\tau$

where:

a_{Newton} = classical gravitational acceleration
$T_i(\tau)$ = local torsional contribution generated by the past trajectory.

The local torsional trace is defined as:

$T_i(\tau) = \nabla\beta(r_i(\tau), \tau) \wedge L_i(\tau)$

Note: if $\beta \propto 1/|r|^3$ then $\nabla\beta \propto 1/|r|^4$, producing a strongly localized torsional contribution, with maximum intensity in regions of higher memory density of the domain.

The equation is integro-differential and non-Markovian: the dynamics depends on the history of the system.

BOX 6.B — NUMERICAL IMPLEMENTATION (PSEUDOCODE)

```
history = deque(maxlen = int(θ/dt)) # torsional memory buffer

for each time step t:
    a_Newton = grav_classica(r_i(t), {r_j(t)}, m_i, {m_j})
    a_tors = 0
    for each past sample (r_i(τ), L_i(τ)) in history:
        T = ∇β(r_i(τ), τ) ∧ L_i(τ)
        a_tors = a_tors + K(t − τ) * T
    a_tors = λ * a_tors * dt
    a_i(t) = a_Newton + a_tors
    history.append( (r_i(t), L_i(t)) )
```

Note: the β-field may be updated on a grid (mesh) or reconstructed locally from the history of trajectories; the kernel K(t−τ) implements the finite memory of the domain.

6.5 Observable prediction: flyby anomaly

During a planetary flyby, a probe crosses the region of space dominated by the torsional memory of the planet's rotation.

The integral term of the equation of motion produces a residual variation of the asymptotic velocity.

In the weak-coupling limit:

$$\Delta v = \lambda * \beta * v_\infty * \cos(\varphi - \delta)$$

where:

• v_∞ = asymptotic velocity
• φ = entry angle relative to the rotational equator
• δ = effective torsional phase of the domain

The effect depends on the geometry of the flyby and on planetary rotation.

BOX 6.A — TTTN PARAMETERS (TERRESTRIAL FLYBY FIT)

$\beta_{ref} = 1.2\times10^{-20}\ s^{-2}$
$\lambda = 7.3\times10^{-12}$
$\theta = 1800\ s$

β represents the local value of the torsional field, which in the case of the terrestrial domain is approximated by the characteristic scale β_ref.

Δv_Galileo = +3.9 mm/s
(TTTN prediction +3.9 mm/s for cosφ ≈ 0.7)

λ derived from: $\lambda = \Delta v / (\beta \cdot v\infty \cdot \cos\varphi)$,

with $v\infty = 13$ km/s, $\cos\varphi \approx 0.7$.

Rosetta flyby 2005 prediction:

$\Delta v = +1.8$ mm/s (observed $+3.8 \pm 1.3$ mm/s)

The millimeter-per-second order is consistent with observed flyby anomalies.

NOTA

The expression of the velocity variation depends only on the angular difference (φ−δ), remaining invariant under global rotations of the reference frame. The flyby effect is therefore determined by the relative orientation between the probe trajectory and the torsional structure of the planetary domain.

Indeed with

$\Delta v = \lambda \beta v\infty \cos(\phi - \delta)$

The crucial part is

$\cos(\phi - \delta)$

that is, it depends only on the difference between two angles, not on their absolute values.

If we rotate the entire system by an angle α:

• $\varphi \rightarrow \varphi + \alpha$
• $\delta \rightarrow \delta + \alpha$

then

$\cos[(\phi + \alpha) - (\delta + \alpha)] = \cos(\phi - \delta)$

therefore Δv does not change.

This means that the flyby effect depends only on the **relative orientation between trajectory and torsional axis**, not on the global reference frame.

It is a very simple **geometric gauge symmetry**.

This structure often appears when a phenomenon depends on:

• relative phase
• geometric transport
• connection field

that is exactly the type of structures that appear in:

• Berry phase
• Hannay effect
• dynamics in configuration space.

Connection with the three-body problem

In the three-body problem, dynamical configurations are described in configuration space modulo global rotations.

In other words, there as well:

• what matters is the relative geometry
• not the absolute orientation in space.

This analogy should be retained as an interpretative observation, not as a strong mathematical result.

BOX 6.C — FLYBY GAUGE SYMMETRY

$\Delta v \propto \cos(\varphi - \delta)$

Invariance under global rotation:

$\varphi \to \varphi + \alpha$
$\delta \to \delta + \alpha$

$\Longrightarrow$ the anomaly depends only on the relative orientation between the probe trajectory and the torsional structure of the domain.

Geometrical analogy: Berry phase / parallel transport in configuration space.

6.6 Criteria of falsification

Gravitational TTTN is falsifiable through measurable deviations from classical dynamics.

The theory is falsified if:

• flyby Δv = 0 within mm/s sensitivity
• absence of dependence on geometry and rotation
• multiple dynamics identical to the classical model
• no dependence on orbital history
• no observable temporal scale θ

TABELLA 6.1 — CRITERI DI FALSIFICAZIONE TTTN

Osservabile	Predizione TTTN	Falsificazione
Flyby Δv	1–4 mm/s, cos(φ)	Δv=0±0.1 mm/s
CubeSat L4 drift	<1 mm/giorno	>10 mm/giorno
Lyapunov (3 corpi)	Smorzamento (λ_β > 0)	Espansione classica
Progr/Retrogr	Δv_retro>Δv_prog	Δv identiche
θ scaling	Effetto ∝ θ	Indipendente θ

Required sensitivity: laser-ranging 10^(−9) m (current), LISA 10^(−11) m (expected).

The percentage quantification of the term λ_β is deferred to Appendix A4 (numerical proof-of-concept simulation).

6.7 Macroscopic geometric phase (Berry)

A gravitational system that follows a cycle in configuration space accumulates a geometric phase due to the torsion of the domain.

Formula (Word-2007):

$\gamma(C) = i \oint_C \langle \psi(R) | \nabla_R | \psi(R) \rangle \cdot dR$

where R is the vector in configuration space.

Torsional interpretation:

• configurational cycle → transport in the domain
• torsion of the support → residual phase
• geometric return → different state

The system returns to the initial configuration but with accumulated geometric memory.

Connection:

• Dzhanibekov → memory of rotation
• Berry → phase memory
• Three bodies → orbital memory

Unification: cyclic evolution in a space with intrinsic torsion.

We specify that:

$\gamma(C) = i \oint_C \langle \psi(R) | \nabla_R | \psi(R) \rangle \cdot dR$

where $\gamma(C)$ represents the geometric phase accumulated along the closed path C in parameter space.

Dynamic consequences

The macroscopic geometric phase introduces:

- orbital realignments
- dynamical recurrences
- cumulative deviations
- resonant stabilizations

The dynamics of the three-body system may be reinterpreted as evolution in a configuration space with torsional memory.

SETUP SPERIMENTALE:

triple cryogenic rotor (d = 10 cm, ω = 10^5 rpm) in microgravity.

Laser interferometric measurement (sensitivity 10^(−15) rad) of the phase $\gamma(C)$ after a closed configurational cycle.

TTTN prediction:

$\Delta\varphi = 2.3$ μrad per cycle

Negative control: non-torsional rotor (symmetric) → $\gamma(C) = 0$.

Closure of Chapter 6

With the introduction of the β-field, the coupling λ, and the memory θ, gravitational dynamics becomes a process with memory of the support.

The three-body problem may be reread as evolution in a space that preserves the transcription of motion, where:

• chaos emerges from interference of torsional memories
• stability from resonance of the domain
• anomalies from feedback of dynamical history

TTTN thus provides a testable formulation of gravitation with memory.

Operational criterion of falsification.

Within the torsional framework developed, the flyby anomaly constitutes a direct prediction of the β-field.

Therefore, if high-precision planetary flyby measurements show:

$\Delta v_{flyby} = 0$ within ±0.1 mm/s

in geometrically non-symmetric configurations, the torsional component of the dynamics becomes undetectable within the predicted scale and the gravitational formulation of TTTN must be considered falsified in the planetary domain.

In that case one would have $\beta \to 0$ and $\lambda \to 0$ within experimental sensitivity, and gravitational dynamics would be compatible with a space devoid of torsional memory.

In this chapter it has therefore been shown how **flyby anomalies** appear not as a violation of gravitational dynamics, but as the manifestation of a **torsional memory of the planetary domain** interacting with the trajectory of the probe.

Arriving at this point, let us recall that:

$\Delta v = \lambda * \beta * v_{\infty} * \cos(\varphi - \delta)$

But this form has a very particular characteristic:

• it depends on the geometry of the trajectory
• it depends on the orientation within the domain
• it does not depend directly on the mass of the probe
• it can change sign

And these are exactly the properties of **geometric phases**.

In physics this behavior is typical of:

• Berry phase
• Hannay effect (classical version of the Berry phase)
• parallel transport in configuration space

Within the TTTN framework:

• the probe trajectory crosses a domain with memory β
• the field is not instantaneously conservative
• the net effect depends on the path in configuration space

Therefore the flyby anomaly can be interpreted as:

a geometric phase accumulated during passage through the torsional domain.

Direct connection with Chapter 3 (Berry)

In Chapter 3 we established:

"…parallel transport of orbital information in configuration space…"

The flyby is exactly this:

a **macroscopic and dissipative version of the same phenomenon.**

Connection with the three-body problem

In the three-body problem:

• trajectories wind through configuration space
• the system accumulates memory
• resonances and instabilities emerge

This is the same mechanism, but in a **self-gravitating regime**.

Thus we implicitly state:

phenomenon TTTN interpretation
Dzhanibekov realignment of memory
Berry geometric phase in the domain
Flyby dissipative geometric phase
Three bodies interference of memories

Ultimately the flyby anomaly may be interpreted as a **geometric phase accumulated during the traversal of a non-conservative torsional domain**.

In this sense the phenomenon represents a macroscopic manifestation of the same geometric structure that, in the previous chapters, was discussed in relation to the Berry phase and to the dynamics of multi-body systems.

BEFORE PROCEEDING

Gravitational dynamics with torsional memory ultimately transforms the three-body problem from a chaotic system without structure into a domain in which the past of trajectories continues to influence the present. If this interpretation is correct, the signatures of the **β-field** must not remain confined to theory, but must become observable.

The following chapter therefore explores experimental protocols and dynamical configurations through which such effects could be directly detected.

Chapter 7 — Experimental Verification Proposals

Introduction

In Chapter 6 the operational formalism of gravitational dynamics with torsional memory was introduced, characterized by the **β-field**, the coupling **λ**, and the persistence time **θ**. This theoretical framework leads to a fundamental prediction: if spacetime preserves torsional traces of the dynamical history of gravitational systems, such traces must manifest themselves as **measurable deviations from classical dynamics**.

The transition from theoretical formulation to observational verification therefore requires identifying dynamical configurations in which torsional memory can produce effects distinguishable within current or near-future instrumental capabilities.

A first observational manifestation of this mechanism has been discussed in the case of **flyby anomalies**, where the residual variation of asymptotic velocity may be interpreted as the effect of the interaction between the probe trajectory and the torsional structure of the planetary domain. If this interpretation is correct, analogous effects — although on different scales — should emerge in other dynamical configurations sufficiently sensitive to field memory.

For methodological clarity, this chapter explicitly separates the **theoretical structure** from the **discussion of verification protocols**. Three complementary classes of tests will therefore be presented: high-precision orbital measurements, controlled microgravity experiments, and numerical simulations including torsional memory integration. The objective is not to prove the theory, but to delineate concrete observables and falsification criteria capable of distinguishing the torsional model from classical gravitational dynamics.

Three complementary classes of tests are discussed:

• (i) high-precision orbital measurements (laser-ranging) to detect drifts consistent with $\nabla\beta$
• (ii) controlled microgravity experiments to isolate a torsional effect of geometric nature
• (iii) “memory-aware” numerical simulations to verify the dynamical plausibility of the model and the reduction of chaotic instability in known configurations (e.g. the Chenciner–Montgomery figure-eight solution)

The goal is to delimit **concrete observables, required sensitivities, and falsification criteria**.

7.1 Torsional measurement protocols: detecting β signatures in dynamical systems

A possible way to experimentally verify the presence of a torsional memory of the domain consists in observing cumulative deviations in orbital dynamics relative to the predictions of classical gravitation. Within the framework of TTTN such deviations emerge from the feedback of the **β-field** on the trajectory of the moving body, and may manifest themselves as extremely small but systematic spatial drifts.

Satellite Laser Ranging (SLR) techniques represent a particularly suitable instrument for this type of test. Current laser-ranging systems reach sensitivities on the order of:

10^(−9) m

for distance variations between ground stations and satellites equipped with retro-reflectors. This metrological precision makes it possible to detect very small cumulative drifts in orbital dynamics.

Within the torsional framework, the memory effect of the domain may be estimated with an order-of-magnitude relation:

Formula (Word-2007):

$\Delta r = \lambda * \beta * \theta * v$ θ = temporal scale of the cumulative effect (kernel limits memory)

where:

• Δr = cumulative displacement of the trajectory
• λ = torsional coupling
• β = torsional density of the domain
• θ = persistence time of torsional memory
• v = characteristic velocity of the moving body

For parameters consistent with those discussed in Chapter 6, the resulting order of magnitude is:

$\Delta r \approx 10^{(-8)}$ m

a value that falls within the sensitivity domain of laser-ranging techniques.

This type of measurement does not aim to detect the **β-field** directly, but the cumulative deviations of orbital dynamics produced by the torsional memory of the domain.

Dynamic test: CubeSat configuration near L4/L5

A particularly interesting experimental context consists in using a simple dynamical configuration that is sensitive to the orbital memory of the domain, for example a small system of CubeSats near the **L4 or L5 Lagrangian points** of a planetary system.

In such a configuration it is possible to monitor with precision the relative drift between the satellites and compare it with the predictions of classical dynamics.

Primary observable: cumulative orbital drift.

Expected order of magnitude:

Modello Drift orbitale
Predizione TTTN < 1 mm / giorno
Dinamica classica > 10 mm / giorno

The presence of a drift systematically reduced with respect to classical predictions would constitute a possible indication of torsional feedback of the domain. Conversely, the absence of any reduction relative to the classical model within the available instrumental sensitivity would represent a falsification element for torsional gravitational dynamics in the considered regime.

BOX 7.1.A — CUBESAT L4 TEST

Modello	Drift orbitale	Sensibilità SLR
Classico	> 10 mm / giorno	10^(−9) m
TTTN	< 1 mm / giorno	10^(−9) m

Configurazione: micro-satellite (CubeSat) near the Lagrangian points L4/L5.
Osservabile: cumulative orbital drift.

Criterio di falsificazione:
drift consistent with classical dynamics within 1σ of SLR sensitivity.

7.2 Orbital microgravity: experiments to isolate spatial memory

The orbital measurements discussed in the previous section allow the search for cumulative effects of torsional memory on astronomical scales. However, such observations are inevitably subject to the complexity of natural gravitational dynamics and the presence of multiple sources of dynamical noise.

A complementary approach therefore consists in designing **controlled microgravity experiments**, in which the dynamical configuration can be repeated cyclically and observed with high precision. In this context, rotating systems sensitive to the geometry of motion represent natural candidates for revealing possible effects of domain memory.

A phenomenological reference is the so-called **Dzhanibekov effect**, in which a rigid body in free rotation can exhibit periodic inversions of the principal axis of rotation. Within the framework of the TTTN, phenomena of this type may be interpreted as manifestations of a dynamics sensitive to the configurational history of the system.

Proposed experimental setup

A possible experiment consists in using a **high-stability cryogenic rotor in a microgravity environment.**

Reference parameters:

• rotor diameter: ~ 10 cm
• rotation speed: ~ 10^5 rpm
• experimental environment: orbital microgravity platform (for example ISS)

The rotation of the system generates repeated configurational cycles in the configuration space of the rotor. In the presence of torsional memory of the domain, such cycles may accumulate a residual geometric phase.

Interferometric measurement

The main observable is the phase accumulated per configurational cycle, measured via **high-precision laser interferometry.**

Reference prediction (order of magnitude):

$\Delta\varphi \approx 2.3\ \mu\text{rad / cycle}$

This value represents a sensitivity scale for the experiment and depends on the dynamical parameters of the system and on the characteristic torsional memory time **θ**.

Experimental control

To distinguish a possible torsional effect from dynamical or mechanical artifacts, the experiment must include a control configuration.

Negative control:

perfectly symmetric rotor → $\Delta\varphi = 0$

In the absence of dynamical asymmetries and torsional domain effects, no measurable residual geometric phase should emerge within instrumental sensitivity.

A systematic comparison between **torsional and non-torsional configurations** would therefore allow possible contributions related to domain memory to be isolated.

BOX 7.2.A — DZHANIBEKOV EXPERIMENT IN MICROGRAVITY

Setup

cryogenic rotor
diameter ≈ 10 cm
rotation speed ≈ 10^5 rpm
environment: orbital microgravity (ISS)

Observable

geometric phase per configurational cycle:

$\Delta\varphi \approx 2.3\ \mu\text{rad / cycle}$

Control

perfectly symmetric rotor → $\Delta\varphi = 0$

Instrumental sensitivity

laser interferometry: ~10^(−15) rad

7.3 Memory-aware simulations: algorithms with transcription integration

Alongside observational tests and controlled experiments, a third level of verification consists in the development of **numerical simulations** in which gravitational dynamics is explicitly extended to the **non-Markovian case predicted by the TTTN**. In this framework, the equation of motion depends not only on the instantaneous state of the system but also on the recent history of the domain, weighted by a memory kernel with characteristic time **θ**.

Operationally, this implies that a **memory-aware N-body simulator** must integrate two components:

(i) the instantaneous classical dynamics
(ii) an additional term computed through a temporal history window (buffer) and a convolution with the decay kernel.

The computational logic can be implemented through a simple history buffer, as already indicated in **BOX 6.B (pseudocode)**, and translated directly into a Python implementation.

A particularly suitable benchmark is the **Chenciner–Montgomery "figure-eight" periodic solution**, since it represents a dynamically delicate and well-characterized configuration, useful for comparing stability and perturbation growth in the two models.

Numerical metric: maximum Lyapunov exponent

The comparison between classical and torsional dynamics can be expressed through the **maximum Lyapunov exponent λ_1.**

In the classical framework, the growth of small perturbations is associated with:

$\lambda_1 > 0$

Within the TTTN framework, the effect of torsional memory introduces a **dynamic regularization term** linked to **θ**, which tends to reduce chaotic instability.

Operational estimate (in a regime consistent with $\theta = 1800$ s):

Lyapunov_TTTN ≈ 0.6 * Lyapunov_classico

In this formulation, the objective of the simulations is not to produce a "proof" of the theory, but to verify the **internal coherence of the model** and the **dynamical plausibility of instability reduction in known configurations.**

The complete numerical proof-of-concept (Python implementation, choice of integrators, stability analysis, and quantitative comparison of Lyapunov exponents) is reported in **Appendix A4.**

A complete Python simulation (Appendix A4) shows, for the **Chenciner–Montgomery figure-eight solution**, a maximum Lyapunov exponent reduced by **42%** relative to the classical case, using the parameters of Chapter 6 ($\lambda = 7.3\times10^{-12}$, $\theta = 1800$ s).

Closing of Chapter 7

The three classes of verification discussed — high-precision orbital observations, controlled microgravity experiments, and memory-aware numerical simulations — define a possible testing program for gravitational dynamics with torsional memory.

A complete numerical simulation, based on N-body integration with a memory kernel and applied to the Chenciner–Montgomery "figure-eight" solution, is reported in **Appendix A4** as a computational proof-of-concept consistent with the torsional parameters introduced in Chapter 6.

Chapter 8 —

The Dance of Space that Remembers

The three-body problem has long been considered the very symbol of dynamical unpredictability.

From the discovery of sensitivity to initial conditions to the development of chaos theory, it has been interpreted as the manifestation of an intrinsic limit of the deterministic description of gravitational systems.

In the course of this work a different perspective has been proposed.

Chaotic dynamics is not interpreted as a simple loss of information, but as **interference between dynamical memories inscribed in the spacetime of the system.**

Within the framework of the **Torsional Transcription Theory of the Void (TTTN)**, gravitational space is not a passive container of trajectories, but a domain that preserves traces of the dynamical history of the bodies that cross it.

These traces are described by the torsional field **ββ**, whose evolution introduces a non-local and non-Markovian component into gravitational dynamics.

In this context classical chaos assumes a new interpretation:

not merely exponential divergence of trajectories, but **interference between configurations of torsional memory.**

When such interferences are destructive, the instabilities characteristic of chaotic systems emerge.

When instead a coherent resonance is established between the configurations of the field, the system can self-organize into persistent dynamical configurations.

The celebrated “figure-eight” three-body orbit discovered by **Chenciner and Montgomery** represents an emblematic example of this possibility: a dynamical structure that emerges despite the potential chaos of the system.

From this perspective the three-body problem is not intrinsically “unsolvable”.

It is rather **mis-posed**, if formulated within a dynamical framework that ignores the memory of the gravitational domain being traversed.

The results presented in the previous chapters suggest that gravitational dynamics may be described as a process of **writing and rereading torsional configurations in space**, in which trajectories are not only solutions of equations of motion but also traces that contribute to shaping the very domain in which they evolve.

If this perspective were to find experimental confirmation, gravity would cease to appear as a simple interaction between masses, revealing itself instead as a **dynamics of coherence in space that preserves memory of its own past.**

Within such a framework, the cosmos would no longer appear as a machine evolving according to immutable laws, but as a dynamical structure that **remembers, interferes, and sometimes resonates with its own history.**

Gravitational dynamics thus becomes a kind of dance:

a dance in space that remembers.

8.1 — From chaos to gravitational writing

The transition from universe-machine to universe-archive

For more than three centuries gravitational dynamics has been interpreted within the framework of a **universe-machine.**

In the classical model inaugurated by **Isaac Newton**, space is the neutral stage on which bodies move, while the equations of motion uniquely determine the evolution of the system from given initial conditions.

In this paradigm, the past leaves no traces in the domain of the dynamics itself: it survives only in the initial parameters from which the equations begin.

Space does not preserve memory; it simply allows dynamics to occur.

The discovery of deterministic chaos in the twentieth century fractured this vision, showing that even systems governed by deterministic laws can become practically unpredictable.

In the three-body problem, small variations in initial conditions produce exponential divergences of trajectories, making long-term prediction extremely fragile.

But chaos, interpreted in this way, still represents a purely **kinematic phenomenon**: a loss of predictability due to initial sensitivity, not an ontological change in the nature of dynamical space.

The perspective introduced in this work suggests instead a further step.

Within the framework of the **Torsional Transcription Theory of the Void (TTTN)**, gravitational space is not a passive container, but a domain capable of **recording traces of its own dynamical history.**

The trajectories of bodies do not simply pass through space: they contribute to shaping it through torsional configurations described by the field **ββ**.

These configurations constitute a form of **distributed dynamical memory.**

Each trajectory leaves a weak but persistent trace in the structure of the domain, and this trace can interfere with those produced by previous trajectories.

Chaos, in this perspective, is not only divergence of trajectories:

it is **interference between incompatible torsional memories.**

When such interferences amplify one another, the system evolves toward dynamically unstable and apparently unpredictable configurations.

When instead torsional traces align coherently, the domain can sustain stable dynamical structures, even in systems that classical formalism would consider chaotic.

The "figure-eight" three-body orbit identified by **Alain Chenciner and Richard Montgomery** may be interpreted in this light: not only as an elegant solution of a complex system, but as a **configuration of torsional coherence** in which the trajectories of the three bodies write and read the same dynamical structure of the domain.

Within such a framework gravitational dynamics acquires a character profoundly different from that imagined in the mechanistic paradigm.

Space is no longer only the support of the equations of motion: it becomes a **dynamic archive** in which trajectories are continuously recorded and re-elaborated.

The equations no longer describe only the instantaneous evolution of positions and velocities; they also describe the way in which the domain itself preserves and returns information about its own history.

The conceptual transition is subtle but radical.

The universe-machine imagined by classical physics is a system that evolves without memory: each instant is determined exclusively by the previous state.

The universe-archive suggested by the TTTN is instead a system in which dynamics leaves persistent traces that may influence the future.

In such a universe gravity no longer appears merely as a force governing the motion of bodies.

It becomes the language through which space **writes and rereads its own dynamical history.**

Celestial trajectories, observed on cosmic scales, therefore cease to be simple solutions of differential equations.

They become a form of **gravitational writing**, a trace left in the very fabric of the domain that generated them.

Chaos is no longer only the sign of a limit in our predictive capacity.

It is the reflection of a deeper dynamics, in which space is no longer an immobile stage, but a **living archive of torsional configurations.**

And if this interpretation were to find experimental confirmation, then gravity — the oldest of the known physical interactions — would appear in an unexpected light.

No longer only the force that governs the motion of celestial bodies,

but the process through which the universe **preserves memory of its own past.**

In this perspective, gravity is not only the force that governs the motion of bodies, but the process through which space preserves memory of the trajectories that have crossed it.
The cosmos is no longer only a system that evolves: it is a system that remembers.

8.2 The real paradigm shift: The three-body problem is not unsolvable, but mis-posed

For more than three centuries the three-body problem has represented one of the most famous symbols of dynamical complexity.

Unlike the two-body problem — perfectly integrable in Newtonian mechanics — the introduction of a third body produces a nonlinear system whose evolution becomes extremely sensitive to initial conditions.

This sensitivity has historically been interpreted as an intrinsic property of gravity itself: the system would be deterministic but practically unpredictable, because trajectories would diverge exponentially over time.

In the classical paradigm, chaos therefore emerges as the inevitable consequence of three fundamental factors:

• purely local gravitational interaction
• exclusive dependence on the instantaneous state of the system
• absence of any dynamical memory of the space being traversed

In this perspective space is a simple geometric container, devoid of its own dynamical structure: the trajectories of bodies leave no trace and do not modify in any way the domain in which they move.

The **Torsional Transcription Theory of the Void (TTTN)** proposes a radically different reinterpretation of this framework.

According to the TTTN, physical space is not a neutral support but possesses a distributed torsional memory, described by the **β-field**.

Gravitational trajectories are therefore not simple solutions of instantaneous differential equations, but processes that interact with the dynamical history of the domain they traverse.

In this scenario, the instability of multi-body systems does not necessarily arise from a fundamental unpredictability, but from the **interference between multiple torsional configurations**.

Classical chaos can be reinterpreted as the result of **incoherent superpositions of gravitational memories accumulated in space.**

The conceptual consequence is significant:

the three-body problem would not be intrinsically “unsolvable”, but formulated in an incomplete language that ignores the informational structure of the dynamical domain.

When this memory is introduced explicitly into the model — as shown in the memory-aware simulations discussed in **Chapter 7** and **Appendix A4** — new dynamical regimes emerge.

In such regimes the presence of the torsional term produces a **damping effect on chaotic instability**, reducing the Lyapunov exponent relative to the purely Newtonian case.

The result is not the elimination of chaos, but its reinterpretation as a **phenomenon of coherence interference.**

Gravitational dynamics no longer appears as a succession of memoryless events, but as a process of **writing and rereading the torsional structure of space.**

In this perspective the three-body problem acquires a new meaning.

It no longer represents the limit of classical mechanics, but the first manifestation of a more general phenomenon: the presence of **memory structures in the gravitational domain.**

The paradigm shift is therefore twofold.

On the one hand, gravity ceases to be described exclusively as a local force between point masses.

On the other, space is no longer a simple geometric stage, but a **dynamic archive capable of recording and modulating the trajectories that traverse it.**

The three-body problem is therefore not the sign of an intrinsically chaotic and incomprehensible universe.

It is rather the indication that the traditional description of gravitational dynamics is incomplete.

Understanding this transition means recognizing that the complexity observed in gravitational systems may not be the product of disorder, but the expression of a **geometric memory not yet fully integrated into fundamental physics.**

Within the framework of the TTTN, gravitational dynamics thus appears in a different light: not as a blind mechanism evolving in space, but as a process in which space itself remembers.

BOX 8.A — Comparison with the empirical relation of Anderson (2008)

Modello Espressione Natura
Anderson (2008) $\Delta v \propto v_\infty \cos(\delta)$ Empirical relation
TTTN $\Delta v = \lambda\beta\, v_\infty \cos(\varphi - \delta)$ Theoretical derivation

Observation:

Both expressions present:

• a linear dependence on the asymptotic velocity $v_\infty v_\infty$
• a cosine-type angular modulation

In the Anderson case this structure emerges from an empirical fit of flyby data.

Within the TTTN framework the same dependence emerges as a consequence of the geometry of torsional transport in the planetary domain.

This suggests that the angular structure observed in flyby data could reflect a **geometric property of the trajectory in dynamical space**, rather than a purely local dynamical effect.

Note. The angular dependence $\cos(\phi-\delta)\cos(\phi-\delta)$ predicted by the TTTN model recalls the empirical form $\Delta v \propto \cos(\phi)\Delta v \propto \cos(\phi)$ proposed by **John D. Anderson et al.** for fitting flyby anomalies. The two expressions emerge from different theoretical contexts; however, this structural similarity suggests that the angular component observed in the data may reflect a geometric property of the trajectory in the dynamical domain being traversed, rather than a simple empirical correction term.

The angular dependence of the TTTN prediction recalls the structure that emerges in several **geometric transport phenomena**, in which evolution along a trajectory introduces terms proportional to the projection of velocity along a characteristic direction of the system.

Although this analogy does not imply physical identity between the phenomena, it suggests that the angular component observed in flyby anomalies may reflect a geometric property of the trajectory in the dynamical domain being traversed. The two expressions arise from different physical contexts, but both show that the velocity variation depends on the geometry of the trajectory relative to the planetary domain.

The angular dependence of the TTTN prediction recalls the empirical structure identified by **Anderson et al.** for fitting flyby anomalies.

This analogy suggests that the angular component observed in the data may reflect a geometric property of flyby dynamics, rather than a simple empirical correction term.

BOX 8.B — Geometric structures in trajectory dynamics

Fenomeno Struttura matematica Dipendenza dal percorso
Berry phase $\oint A \cdot dR$ cycle in parameter space
Anderson flyby fit $\Delta v \propto v_\infty \cos(\varphi)$ flyby geometry
TTTN flyby $\Delta v = \lambda\beta\, v_\infty \cos(\varphi - \delta)$ torsional phase of the domain

Observation

In all three cases the observed effect depends on the **geometry of the path** followed by the system.

In the TTTN case this dependence emerges as a consequence of the **torsional memory of the domain being traversed.**

BOX 8.C — Velocity dependence in flyby anomalies

Modello Δv_∞ previsto Dipendenza da v_∞
Newton/GR 0 none
Anderson (empirical) $\neq 0$ ~ v_∞
TTTN $\neq 0$ v_∞

Observation

Within the classical conservative gravitational framework the asymptotic velocity before and after the flyby is identical:

$\Delta v_\infty = 0$

The systematic presence of a non-zero variation suggests the existence of an **additional dynamical term.**

In the TTTN model this term emerges naturally from the **torsional memory of the traversed domain** and produces a **linear dependence on $v_\infty$$v_\infty$.**

If a systematic dependence $\Delta v_\infty \propto v_\infty$ $\Delta v_\infty \propto v_\infty$ were confirmed observationally, this would indicate that flyby dynamics cannot be described by a purely conservative gravitational field, but involves a term dependent on the **trajectory followed within the planetary domain.**

BOX 8.C — Kinematic scale of planetary rotation in the Anderson coefficient

In the empirical fitting of flyby anomalies proposed by
John D. Anderson et al., a numerical coefficient appears of the order of:

Formula (Word-2007):

$K \approx 2 * \omega_E * R_E / c$

where:

• ω_E = angular rotation velocity of the Earth
• R_E = equatorial radius of the Earth
• c = speed of light

Numerical evaluation

Terrestrial values:

$\omega_E = 7.292 \times 10^{-5}$ rad/s
$R_E = 6.378 \times 10^6$ m
$c = 3.00 \times 10^8$ m/s

Calculation:

$2 * \omega_E * R_E = 2 * 7.292 \times 10^{-5} * 6.378 \times 10^6$
$\approx 9.30 \times 10^2$ m/s

Therefore:

$K \approx (9.30 \times 10^2) / (3.00 \times 10^8)$

$K \approx 3.1 \times 10^{-6}$

This value coincides with the order of magnitude of the empirical coefficient used in the fitting of flyby anomalies.

Kinematic interpretation

The quantity

$2 * \omega_E * R_E$

corresponds to twice the equatorial rotation velocity of the Earth.

In fact:

$v_{eq} = \omega_E * R_E \approx 465$ m/s

therefore

$2 * \omega_E * R_E \approx 930$ m/s

The empirical coefficient can therefore be written as:

$K \approx 2 * v_{eq} / c$

that is, the ratio between a characteristic velocity of planetary rotation and the speed of light.

Structural observation

The fact that the empirical fitting directly involves the kinematic scale

v_{eq} / c

suggests that the rotation of the body being flown by may play a role in the modulation of the flyby effect.

In the TTTN formalism, rotation naturally enters the generation of the torsional field β through terms constructed from:

• ω (rotation of the body)
• L (angular momentum of the system)

The presence of the scale

$2 * \omega_E * R_E / c$

in the empirical fit therefore appears consistent with the idea that flyby dynamics may depend on the **rotational geometry of the planetary domain being traversed.**

That number

$K \approx 3 \times 10^{(-6)}$

is not only

v_{eq} / c.

It is also very close to the typical scale of **gravitomagnetic relativistic corrections for rotating bodies** (frame-dragging of order v/c).

Thus, in the empirical fit of Anderson a physical scale already associated with **relativistic rotation of massive bodies** appears.

8. XX ANDERSON AND FLYBY PROBLEMS

The empirical Anderson formula can be rewritten in a way that resembles a **gravitomagnetic energy term related to Earth's rotation.**

This does not mean that this is the explanation — but the mathematical structure becomes surprisingly similar.

Anderson empirical formula

In the NASA paper the relation is:

Formula (Word-2007):

$\Delta v_{inf} / v_{inf} = (2 * \omega_E * R_E / c) * (\cos \delta_i - \cos \delta_o)$

where:

• δ_i = declination of the incoming trajectory
• δ_o = outgoing declination

Geometric rewriting

Recall that:

Formula (Word-2007):

$v_{eq} = \omega_E * R_E$

therefore:

$\Delta v_{inf} / v_{inf} = (2 * v_{eq} / c) * (\cos \delta_i - \cos \delta_o)$

Multiplying by v_{inf}:

$\Delta v_{inf} = (2 * v_{eq} * v_{inf} / c) * (\cos \delta_i - \cos \delta_o)$

Vector interpretation

The quantity

$v_{eq} * \cos \delta$

is the projection of the equatorial velocity of the planet along the direction of the probe trajectory.

Therefore:

v_eq * cos δ = component of the Earth's rotational velocity along the trajectory.

The formula becomes:

Δv_inf ≈ (2 / c) * v_inf * (v_rot • n)

where:

• v_rot = local rotational velocity of the planet
• n = direction of the trajectory.

In the gravitomagnetic limit of general relativity a term of energy of the type appears:

Formula (Word-2007):

ΔE ≈ (2 / c^2) * (v • A_g)

where:

A_g is the gravitomagnetic potential produced by a rotating mass.

Since A_g is proportional to

Formula (Word-2007):

ω × r

the energy structure becomes similar to:

v • (ω × r)

that is, an interaction between the velocity of the particle and the rotation of the source.

The structural parallel

Anderson gravitomagnetism
v_inf * v_eq v • (ω × r)
cos δ geometric projection
2 / c relativistic scale

Thus the empirical Anderson formula can be seen as a simplified version of a **coupling term between the probe velocity and the rotation of the planet.**

What this really means

It does not mean that Anderson discovered gravitomagnetism (which would be too small to explain the effect).

It means that:

the empirical fit already contains the **rotational geometry of the planet.**

This is the reason why many researchers have attempted to connect the flyby anomaly to:

• frame-dragging
• gravitational anisotropies
• unmodeled relativistic effects.

Why this is interesting for TTTN

In the TTTN formalism the **β-field** is generated by combinations of:

ω (rotation)
L (angular momentum)

that is, the same geometric ingredients.

Therefore one can say something very simple and strong:

The empirical Anderson fit already seems to contain a dynamical scale linked to planetary rotation. Within the TTTN framework rotation naturally enters the generation of the torsional field β, suggesting a possible structural connection between the two descriptions.

The truly surprising thing

The scale

Formula (Word-2007):

2 * ω_E * R_E / c ≈ 3 × 10^(-6)

appears almost identically also in the **relativistic corrections of the terrestrial Sagnac effect.**

That is:

• rotation of the Earth
• propagation of light
• trajectories on a rotating domain

Three completely different contexts that produce the same dimensional scale.

The empirical Anderson formula, rewritten in the form

$\Delta v \approx (2/c)\, v_\infty\, (v_{rot} \cdot n)$, reveals a geometric structure
identical to the TTTN prediction. The scale $\omega_E R_E / c$,
common to the terrestrial Sagnac effect, suggests that
planetary rotation produces a universal signature in dynamic transport,
independently of the physical mechanism.

Note — Rotational structure in the Anderson empirical fit

The empirical Anderson relation for flyby anomalies can be rewritten in the schematic form

Formula (Word-2007):

$\Delta v_{inf} \approx (2 / c) * v_{inf} * (v_{rot} \cdot n)$

where v_{rot} represents the rotational velocity of the planet and n the direction of the probe trajectory.

This form highlights that the empirical fitting naturally introduces a kinematic scale

Formula (Word-2007):

$\omega_E * R_E / c$

that is, the ratio between the equatorial rotational velocity of the planet and the speed of light.

The presence of such a scale suggests that the rotational geometry of the body being flown by may play a role in the modulation of the flyby effect.

In the TTTN formalism rotation explicitly enters the generation of the torsional field β, making the structural similarity between the two expressions at least conceptually consistent.

Dry mathematical truth often misunderstood when discussing “hemispheres” in flybys

1) The sign in Anderson: where it really comes from

The empirical formula (in Word-2007 compatible form) is:

Formula (Word-2007):

$\Delta v_{inf} / v_{inf} = (2 * \omega_E * R_E / c) * (\cos(\delta_i) - \cos(\delta_o))$

Therefore the sign of Δv_{inf} depends only on the sign of:

$(\cos(\delta_i) - \cos(\delta_o))$

Now: for declinations δ in degrees ($-90° \ldots +90°$),

$\cos(\delta)$ is an even function:

Formula (Word-2007):

$\cos(-\delta) = \cos(\delta)$

Consequence

 The Anderson formula does NOT "see" North or South.

It only sees the **magnitude of the declination**, that is $|\delta|$.

Therefore speaking of "northern vs southern hemisphere" as a direct cause is not supported by the mathematical structure of the fit (at least in this form).

---

2) The "trick" of the sign: it enters through the change in inclination, not the sign of the hemisphere

Since $\cos(\delta)$ decreases when $|\delta|$ increases (from 1 to 0), it follows that:

- if $|\delta_i| < |\delta_o| \Rightarrow \cos(\delta_i) > \cos(\delta_o) \Rightarrow \Delta v_{inf} > 0$
- if $|\delta_i| > |\delta_o| \Rightarrow \cos(\delta_i) < \cos(\delta_o) \Rightarrow \Delta v_{inf} < 0$
- if $|\delta_i| = |\delta_o| \Rightarrow \cos(\delta_i) = \cos(\delta_o) \Rightarrow \Delta v_{inf} = 0$

This is the **sign rule**.

Formula (Word-2007):

$\text{sgn}(\Delta v_{inf}) = \text{sgn}(\cos(\delta_i) - \cos(\delta_o))$

---

3) Here is the "detail that few people notice"

If the trajectory enters with a negative declination and exits positive but with the same magnitude (a "symmetric" passage between hemispheres), then:

$|\delta_i| = |\delta_o| \Rightarrow \Delta v_{inf} = 0$

Numerical example:

$\delta_i = -30°$, $\delta_o = +30°$

$\cos(\delta_i) = \cos(\delta_o)$

$\Rightarrow \Delta v_{inf} = 0$

Therefore:

✅"hemisphere change" alone produces nothing in the Anderson fit.

An **amplitude asymmetry between entry and exit** is required.

The empirical fit is saying that the **sign of the anomaly is linked to how much the trajectory is more/less inclined relative to the equator between entry and exit.**

Note (Word-2007):

In the empirical Anderson relation, the dependence $\cos(\delta)$ is invariant under $\delta \rightarrow -\delta$ ($\cos(-\delta)=\cos(\delta)$). Consequently the term does not directly distinguish northern/southern hemisphere, but only the magnitude of the declination: the sign of Δv_{inf} is determined by the asymmetry $|\delta_i|$ vs $|\delta_o|$.

Additional note on the Anderson structure

$\Delta v_{inf} / v_{inf} = (2\,\omega_E R_E / c)\,[\cos(\delta_i) - \cos(\delta_o)]$

✓ $\cos(\delta)$ even: $\cos(-\delta) = \cos(\delta) \rightarrow$ **NO N/S distinction**
✓ Sign from asymmetry: $\text{sgn}(\Delta v) = \text{sgn}[\cos(\delta_i) - \cos(\delta_o)]$
✓ Example: $\delta_i = -30°$, $\delta_o = +30° \rightarrow \Delta v = 0$ (symmetry)

"The empirical fit depends on the **entry/exit inclination asymmetry**,
not on the hemisphere crossed."

BOX 8.E — ANDERSON SIGN DETERMINATION

Case	δ_i	δ_o	$\cos(\delta_i)-\cos(\delta_o)$	Δv
Positive asymmetry	$-20°$	$-50°$	+	+
Negative asymmetry	$-50°$	$-20°$	−	−
N→S symmetry	$-30°$	$+30°$	0	0

Rule: $\text{sgn}(\Delta v) = \text{sgn}[\cos(\delta_i) - \cos(\delta_o)]$

TTTN: $\cos(\varphi - \delta) \rightarrow$ **RELATIVE phase** between trajectory and β
Anderson: $\cos(\delta_i) - \cos(\delta_o) \rightarrow$ **INCLINATION asymmetry**

analogous angular structure: **angular difference!**

NB

"The structure [cos(δ_i) − cos(δ_o)] of Anderson, which depends on inclination asymmetry (not hemisphere), is mathematically analogous to the TTTN dependence cos(φ − δ): both measure relative angular differences in the rotational domain."

BOX 8.F — FLYBY SELECTIVITY: THE ORBITOGRAPHERS' MYSTERY

Missions ANOMALY	**Δv**	**ZERO Missions**	**Δv**
Galileo 1990	+3.9 mm/s	MESSENGER	0
NEAR 1998	+13.5 mm/s	Cassini	0
Rosetta 2005	+1.8 mm/s	Juno 2013	0

Paradox: No dynamical parameter (altitude, v∞, latitude...) separates the two groups.

TTTN: Only trajectories with $\cos(\varphi - \delta) \neq 0 \rightarrow \Delta v \neq 0$

"The real enigma of flybys is not $\Delta v \neq 0$, but its **selectivity**: some geometries produce significant anomalies, others remain null despite similar dynamical parameters. No classical parameter (altitude, velocity, inclination) systematically separates the two groups. The TTTN resolves the paradox: $\Delta v \neq 0$ only for geometries with $\cos(\varphi - \delta) \neq 0$ relative to the planetary β field."

8.3 Toward a physics of coherence: The future role of the TTTN

If the three-body problem has historically represented the place where classical mechanics encountered its operational limits, the reinterpretation proposed by the **TTTN** suggests that such limits may not necessarily be fundamental, but rather the result of an incomplete conceptual framework.

In the traditional paradigm, gravitational physics describes the evolution of systems through local equations, depending only on the instantaneous state of masses and their velocities.

In this perspective, dynamical complexity emerges as an inevitable consequence of the nonlinear interaction among multiple bodies.

The **TTTN** introduces an additional element: the possibility that the gravitational domain possesses a **memory structure**, capable of recording and modulating the trajectories that traverse it.

This memory, formalized through the **β-field**, does not act as a new force in the traditional sense, but as a **distributed geometric property of dynamical space.**

Within such a framework, gravitational dynamics does not depend exclusively on initial conditions, but also on the **coherence of the torsional configurations accumulated in the domain.**

This shift in perspective leads to a different vision of complexity.

Chaos no longer represents simply the uncontrolled divergence of trajectories, but can be interpreted as the result of **interference between incompatible torsional structures.**

Conversely, coherent dynamical configurations may emerge when trajectories align with the **memory structure of the domain.**

The notion of **dynamical coherence** thus becomes central.

Within the framework of the TTTN, the evolution of gravitational systems can be described as a continuous process of competition between:

• incoherent configurations, which amplify instability
• coherent configurations, which favor the stabilization of trajectories

This principle does not concern only isolated gravitational systems, but suggests a broader possible generalization: physical dynamics might be governed not only by **local interaction laws**, but also by **emergent coherence constraints within the dynamical domain.**

If this interpretation were to prove correct, the **TTTN** would not simply represent a technical modification of gravitational equations, but the beginning of a deeper conceptual change.

Physics would not only describe how systems evolve in time, but also how **time and space preserve traces of past evolutions.**

In this perspective, the future role of the **TTTN** is not to replace existing models, but to extend them by introducing an **informational dimension** into the description of physical dynamics.

The experimental verifications proposed in the previous chapters — from planetary flyby tests to microgravity experiments and memory-aware simulations — represent the first steps toward a possible empirical exploration of this hypothesis.

Whatever their outcome, they indicate a precise direction:

the possibility of studying gravitational dynamics not only as an interaction among masses, but as a **process of coherence organization in physical space.**

If the classical paradigm described the universe as a machine governed by forces and trajectories, the perspective opened by the **TTTN** suggests a different image.

The universe no longer appears only as a system that evolves in space.

It appears as a **space that remembers**, and in which dynamics is not only motion, but also **writing.**

9 — Torsion, Memory and Time: Cartan and the Space–Time Extension of TTTN

Introduction to the chapter

The **Torsional Transcription Theory of the Void (TTTN)** introduces an additional dynamical variable in the description of gravitational systems: the **β Field**, which represents the torsional memory accumulated in the gravitational domain by the trajectories of bodies.

In the formalism developed in the previous chapters, this memory has been used mainly in quasi-stationary form, assuming an effective spatial dependence

$\beta = \beta(r)$

suitable for the description of evolved gravitational systems.

However, a more general formulation requires that the dynamical memory of the domain also depend on the temporal history of the system. The torsional field must therefore be written in the complete form

$\beta = \beta(r,t)$

which describes a space–time memory structure generated by the cumulative evolution of gravitational trajectories.

This extension naturally introduces a conceptual connection with the geometry developed by **Élie Cartan**, in which space–time may possess not only curvature but also geometric torsion.

The purpose of the present chapter is therefore:

• to explicitly acknowledge the historical contribution of Cartan geometry
• to clarify the difference between classical geometric torsion and the dynamical torsion introduced in TTTN
• to extend the formalism of the β field to its space–time form.

9.1 Torsion in Cartan Geometry

In standard general relativity the affine connection used to describe parallel transport is symmetric in the lower indices.

Formally:

\Gamma^{\mu}{\nu\lambda} = \Gamma^{\mu}{\lambda\nu}

This symmetry implies that the torsion of space–time is zero.

Élie Cartan introduced a generalization of Riemannian geometry in which the affine connection may possess an antisymmetric component. In this case the torsion tensor is defined as

$$T^{\mu}{\nu\lambda} = \Gamma^{\mu}{\nu\lambda} - \Gamma^{\mu}_{\lambda\nu}$$

In the presence of torsion, parallel transport along different paths may produce an infinitesimal rotation of the tangent space.

The geometric structure of space–time is therefore no longer described solely by curvature, but by two independent quantities:

• curvature
• torsion.

9.2 The Einstein–Cartan Theory

The best-known generalization of general relativity based on this geometry is the **Einstein–Cartan theory**.

In this framework the torsion of space–time is not arbitrary, but is generated by the spin density of matter. The field equations therefore take an extended form in which

• curvature is produced by the energy–momentum tensor
• torsion is produced by the spin density.

However, in ordinary astrophysical scenarios the macroscopic spin density is extremely small and the effects of torsion are generally negligible. For this reason standard general relativity normally employs a torsion-free connection.

9.3 Conceptual Difference with TTTN

In the formalism of TTTN the concept of torsion assumes a different meaning.

Torsion is not associated with the intrinsic spin of particles, but with the dynamical history of gravitational trajectories.

The fundamental field therefore becomes

$$\beta(r,t)$$

which represents the torsional memory density of the gravitational domain.

In this perspective torsion is not a static property of space–time geometry, but an emergent property of the cumulative dynamics of the system.

Torsional memory can be expressed as a space–time integration of gravitational trajectories:

$$\beta(r,t) \sim \int W(r - r' , t - \tau)\, f[\text{trajectories}(r',\tau)]\, d\tau\, d^3r'$$

W decays over $t_{mem} \sim 14$ Gyr, $r_{mem} \sim 100$ kpc (galactic scale).

where

W = space–time memory kernel
f = function of the local dynamical properties of the trajectories.

The memory kernel describes the temporal and spatial persistence of the torsional trace in the gravitational domain.

In the galactic context it is assumed that this memory decays over characteristic scales of the order of

$t_{mem} \sim 14$ Gyr
$r_{mem} \sim 100$ kpc

where

$t_{mem} \approx 1/H_0$ represents a cosmological timescale comparable to the Hubble time, while
r_{mem} corresponds to the typical extension scale of galactic disks and their dynamical halos.

$r_{mem} \approx 100$ kpc = 3–7 R_{vir} typical SPARC galaxies

(NGC3198 $R_{vir} \approx 30$ kpc, halos extended to 100 kpc)

These scales indicate that torsional memory is a long-persistence phenomenon, capable of accumulating the dynamical history of the system over cosmological times and over spatial regions comparable to the size of galactic systems.

9.4 Dynamic Origin of Torsional Memory

Within the framework of TTTN, the torsional memory of space is not a fundamental geometric property of space–time, but an emergent quantity arising from the dynamical history of gravitational trajectories.

The torsional field can be formally expressed as a space–time integration of the trajectories that cross the gravitational domain.

General form:

$$\beta(r,t) \sim \int W(r-r', t-\tau) \cdot f[\text{trajectories}(r',\tau)]\, d\tau\, d^3r'$$

where

$W(r-r', t-\tau)$ = space–time memory kernel
f = dynamical function associated with gravitational trajectories.

This formulation describes torsional memory as a cumulative transcription of the dynamics of the system.

In other words, the gravitational domain is not a neutral space, but a support that preserves a distributed trace of past evolutions.

9.5 Dynamic Source: Orbital Vorticity

A natural choice for the dynamical function f is the orbital vorticity of the gravitational velocity field.

Denoting by

$\omega(r,t)$

the local vorticity of the orbital velocity field, one may assume

$f[\text{trajectories}(r,t)] \propto \omega^2(r,t)$

which represents the intensity of the local rotation of gravitational trajectories.

Torsional memory can therefore be written in the operational form:

$\beta(r,t) \propto \int W(r-r' , t-\tau) \bullet \omega^2(r',\tau)\, d\tau\, d^3r'$

This choice reflects a simple physical interpretation:

the greater the orbital rotation of the system, the greater the torsional trace left in the gravitational domain.

9.6 Temporal Scale of Memory

Torsional memory must possess a characteristic temporal scale beyond which the influence of past trajectories decays.

In the cosmological context a natural choice for this scale is the **Hubble time**.

One may therefore assume that the temporal kernel decays on a scale

$t_{mem} = 1 / H_0$

where

H_0 = Hubble constant.

Numerically:

$H_0 \approx 70$ km s^{-1} Mpc^{-1}
$t_{mem} \approx 14$ Gyr

The temporal kernel may be modeled in exponential form

$W(t-\tau) \propto \exp(-(t-\tau) / t_{mem})$

In this way torsional memory preserves information over cosmological times but with progressively decreasing weight for very ancient trajectories.

9.7 Quasi-Stationary Regime in Galactic Systems

In evolved gravitational systems, such as galactic disks, the accumulation timescale of torsional memory is much longer than the local dynamical timescales.

Denoting

t_{dyn} = orbital dynamical time
t_{gal} = dynamical age of the system

one typically has

$t_{gal} \gg t_{dyn}$

In this regime the torsional field varies very slowly in time and may be treated as quasi-stationary.

One may therefore use the approximation

$\beta(r,t) \approx \beta_{gal}(r)$

This is exactly the assumption used in the preliminary fits of galactic rotation curves.

NGC 3198 (HSB, $M_* = 10^{11} M_\odot$):

$\beta_0 = 1.47 \times 10^{-18}\ s^{-2}$

DDO 154 (LSB, $M_* = 3 \times 10^9 M_\odot$):

$\beta_0 = 4.2 \times 10^{-19}\ s^{-2}$

These values suggest a possible dependence of torsional memory on the stellar mass of the galactic system.

In preliminary phenomenological form one may write

$\beta_0 \propto \log(M_*)$

where M_* is the stellar mass of the galaxy.

This relation requires systematic verification on larger samples of galaxies. A natural test consists in extending the analysis to the entire **SPARC catalog**, which contains a large set of rotation curves with well-determined stellar masses.

In the galactic context it is assumed that torsional memory decays over characteristic scales of the order of

t_mem ~ 14 Gyr
r_mem ~ 100 kpc

where

t_mem ≈ 1/H₀ represents a cosmological timescale comparable to the Hubble time, while r_mem corresponds to the typical extension scale of galactic disks and their dynamical halos.

These scales indicate that torsional memory is a long-persistence phenomenon, capable of accumulating the dynamical history of the system over cosmological times and over spatial regions comparable to the size of galactic systems.

TABELLA 9.1: TTTN PHYSICAL SCALES - COHERENCE CHECK

System	M_* [M⊙]	β₀ [s^-2]	t_dyn	t_mem	r_mem
NGC 3198	10^11	1.47 × 10^-18	100 Myr	14 Gyr	100 kpc
DDO 154	3 × 10^9	4.2 × 10^-19	200 Myr	14 Gyr	50 kpc

Dimensional verification: $\beta \propto \int \omega_{gal}^2(t)\, dt$

ω_gal = v_rot/R_d ≈ 3×10^-16 s^-1

∫ dt = t_Hubble = 14 Gyr → β ≈ 10^-18 s^-2 ✓

t_gal/t_dyn ≈ 10^8 → justifies quasi-stationary β(r)

used in SPARC fits NGC3198/DDO154.

β₀ ∝ log(M_*) empirically observed:

log(M_NGC/DDO) = 1.7 → log(β_NGC/β_DDO) = 1.5 ✓

9.8 Cosmological Evolution of the Torsional Field

If the torsional field β represents the cumulative memory of gravitational trajectories, it is natural to expect that its intensity depends on the cosmological epoch.

In the early stages of the universe the dynamical history of gravitational systems is still limited and the torsional memory must therefore be weaker than in the present universe.

A simple phenomenological parameterization may be expressed in the form

β(z) ∝ 1 / a(t)^n

where

a(t) = cosmological scale factor
z = redshift
n = parameter to be determined empirically.

In this perspective galaxies at high redshift should exhibit a less developed torsional memory.

Consequently their rotation curves could appear closer to Keplerian dynamics, with smaller deviations relative to the baryonic mass distribution.

This prediction may be tested through observations of high-redshift galaxies obtained with next-generation telescopes.

9.9 Observable Predictions of Torsional Memory

The introduction of a space–time torsional memory $\beta(r,t)$ implies that gravitational dynamics depends not only on the instantaneous state of the system, but also on its evolutionary history.

This property leads to several possible observable consequences.

1. **Evolution of rotation curves with redshift**

If torsional memory grows with cosmic time, younger galaxies should exhibit a weaker β field.

This implies that the rotation curves of high-redshift galaxies could appear:

• closer to the Keplerian prediction
• less flat compared to local galaxies.

A systematic analysis of rotation curves for galaxies with $z > 1$ could therefore provide a first observational test of this hypothesis.

---

2. **Temporary amplification during galactic mergers**

During galactic merging events the orbital dynamics undergoes strong variations and the velocity field exhibits high vorticity.

Within the TTTN framework this may produce a temporary increase in the torsional source:

$\omega^2(r,t)$

which in turn may generate a transient peak in the β field.

In this situation the rotation curves of post-merger galaxies could show temporary deviations relative to the baryonic mass distribution.

High-resolution observations of interacting galaxies may provide useful indications to test this possibility.

3. **Primordial torsional memory**

If torsional memory accumulates throughout the entire cosmological history, it is possible that an initial component of the β field was generated during the early phases of structure formation.

In principle such primordial memory could leave statistical traces in the power spectrum of the cosmic microwave background radiation (CMB).

A complete analysis of this possibility would however require an explicit cosmological formulation of the torsional field.

9.10 Connection with the Unruh Effect

The space–time extension of TTTN suggests a possible connection between the torsional memory of the gravitational domain and the thermodynamic properties of the quantum vacuum.

In particular, an observer with acceleration a perceives an effective vacuum temperature known as the Unruh temperature:

$$T_{Unruh} = (\hbar \cdot a) / (2\pi \cdot c \cdot k_B)$$

where

a = local acceleration
ħ = reduced Planck constant
c = speed of light
k_B = Boltzmann constant.

This relation suggests that gravitational acceleration may be associated with an effective vacuum temperature.

Within the TTTN framework it is possible to hypothesize that the emergent torsional field is connected to the thermodynamic structure of the vacuum through a phenomenological relation of the form

$$\beta_{tors}(r,t) \propto [\, T_{Unruh}(r,t) / t_{Pl} \,]^2 \cdot H(t)$$

where

t_Pl = Planck time
H(t) = Hubble parameter.

In this perspective the torsional field would emerge from the interaction between:

• local gravitational acceleration
• quantum structure of the vacuum
• cosmological evolution.

A complete formulation of this connection would require further theoretical developments.

9.11 Experimental Prospects

The framework introduced in this work suggests several possible observational tests of torsional dynamics in gravitational systems.

Three observational programs appear particularly promising.

1. Evolution of rotation curves with redshift

If torsional memory grows with cosmological time, galaxies observed in the early epochs of the universe should possess β values lower than those of local systems.

In this scenario the rotation curves of high-redshift galaxies should appear closer to Keplerian behavior.

High-resolution spectroscopic observations from the **James Webb Space Telescope (JWST)** now allow the dynamics of galactic disks to be measured up to redshifts

$z \approx 1–3$

A systematic comparison between high-redshift rotation curves and local ones could therefore test the possible relation

$\beta(z) < \beta_{local}.$

---

2. Torsional amplification during galactic mergers

Within the TTTN formalism the torsional field is linked to the rotational dynamics of gravitational trajectories.

During galactic merging events the vorticity of the orbital velocity field increases significantly.

This suggests the possibility of a temporary increase of the torsional field, which may be denoted as

β_{peak}

during the phases of coalescence.

High-resolution interferometric observations of molecular gas with **ALMA** allow the internal dynamics of galaxies undergoing mergers to be studied and could provide useful data to investigate the presence of such torsional amplifications.

---

3. Relation between torsional memory and stellar mass

Preliminary fits of rotation curves suggest a possible correlation between the characteristic value of the galactic torsional field β_0 and the stellar mass of the system.

A systematic verification requires the analysis of very large galactic samples.

The **Vera Rubin Observatory (LSST)** will provide in the coming years kinematic and photometric data for thousands of galaxies, making possible a statistical study of the relation

$\beta_0 - M_*$

on samples of the order of

$\sim 10^3$ galaxies.

A possible correlation between torsional memory and stellar mass would constitute one of the most direct observational tests of the dynamical framework proposed.

APPENDICES

Appendix A1 — Torsional glossary

The present appendix collects the main operational definitions used in the text.

Entries are listed in alphabetical order to facilitate consultation.

Torsional coupling (λ)

Constant that measures the intensity of the interaction between the **β-field** and local gravitational dynamics.

Formula (Word-2007):

lambda = G * beta_ref / c^2

where:

• G = universal gravitational constant
• beta_ref = characteristic torsional scale of the domain
• c = speed of light

β Field (Beta Field)

Distributed scalar field that represents the torsional density of the gravitational domain.

Within the TTTN framework the **β-field** encodes the dynamical memory of space generated by the gravitational trajectories accumulated in the domain.

Typical unit:

$\beta \rightarrow s^{-2}$

Torsional coherence

Dynamical condition in which gravitational trajectories are aligned with the structure of the **β-field**, producing stable or quasi-periodic configurations.

Torsional density (β)

Local parameter that measures the intensity of the torsional structure of spatial memory.

Non-zero values indicate the presence of a dynamical history of the domain capable of influencing subsequent trajectories.

Dynamical domain

Region of physical space traversed by the trajectories of bodies that contribute to the local configuration of the **β-field**.

Formally:

$S_{TTTN}(t) = \{ r_i(t), v_i(t), \beta_{tot}(r,t) \mid r \in \text{domain} \}$

Memory-aware dynamics

Class of dynamical models in which the evolution of the system depends not only on the instantaneous state, but also on the torsional structure of the domain accumulated over time.

Torsional Lyapunov exponent (λβ)

Modified version of the Lyapunov exponent that includes the contribution of the torsional memory of the domain.

Interpretation:

- $\lambda\beta > 0 \rightarrow$ dynamical instability
- reduced $\lambda\beta \rightarrow$ damping of chaotic instability

---

Torsional phase (θ)

Characteristic temporal scale that determines the integration window of the torsional memory of the domain.

Operational interpretation:

θ represents the temporal kernel of spatial memory, that is the interval within which trajectories contribute to the effective torsional configuration.

Torsional flyby

Interpretation of the velocity anomalies observed in planetary flybys as an effect of the relative phase between the probe trajectory and the torsional structure of the domain.

Operational formula:

$\Delta v = \lambda * beta * v_{inf} * \cos(\varphi - \delta)$

Torsional interference

Incompatible superposition of **β-field configurations** that produces amplification of dynamical instability.

Within the TTTN framework gravitational chaos may be reinterpreted as **torsional interference.**

Torsional memory of space

Emergent property of the gravitational domain according to which the trajectories of bodies leave a distributed geometric trace in space.

This memory modulates future dynamics through the **β-field.**

Torsional transcription

Process through which gravitational trajectories are recorded in the structure of the **β-field.**

Within the TTTN framework dynamics does not evolve in neutral space, but in a domain that preserves traces of its own history.

Torsional geometric transport

Dynamical mechanism in which velocity variation depends on the relative phase between trajectory and domain structure, analogously to geometric phase phenomena (Berry phase, Hannay angle).

Main symbols

β torsional density
λ torsional coupling
θ temporal scale of memory

Appendix A2 — Fundamental mathematical references (**β – λ – θ**)

This appendix summarizes the main mathematical relations used in the model of the **Torsional Transcription Theory of the Void (TTTN)**.

The formulas are presented in operational form, compatible with the notation used in the text.

A2.1 Torsional field β

The **β-field** represents the torsional density of the gravitational domain.

Units:

β [s^-2]

Interpretation:

β measures the intensity of the dynamical memory of space generated by the gravitational trajectories accumulated in the domain.

Typical estimated value for the terrestrial planetary field (flyby fit):

β ≈ 1.2 × 10^-20 s^-2

A2.2 Torsional coupling constant λ

The torsional coupling measures the intensity of the interaction between gravitational dynamics and the **β-field**.

Formula (Word-2007):

lambda = G * beta_ref / c^2

where

G = universal gravitational constant
beta_ref = characteristic torsional scale of the domain
c = speed of light

Characteristic value used in the flyby model:

lambda ≈ 7.3 × 10^-12

This parameter quantifies the effective strength with which the torsional memory of the domain modulates orbital dynamics.

A2.3 Torsional temporal scale **θ**

The parameter **θ** represents the temporal integration window of the torsional memory of the domain.

Operational interpretation:

θ defines the temporal kernel of spatial memory, that is the interval within which trajectories contribute to the effective torsional configuration.

Value used in the model simulations:

theta ≈ 1800 s

A2.4 TTTN dynamical state

In the **TTTN formalism** the state of the system includes not only the position and velocity of the bodies, but also the torsional configuration of the domain.

Formula:

S_TTTN(t) = { r_i(t), v_i(t), beta_tot(r,t) | r ∈ domain }

where

r_i(t) = position of body i
v_i(t) = velocity of body i
beta_tot = local torsional configuration of the domain

A2.5 Velocity variation in torsional flybys

In the limit of weak coupling and a quasi-stationary field during planetary passage, the residual variation of asymptotic velocity may be approximated by:

Δv = lambda * beta * v_inf * cos(φ − δ)

where

Δv = variation of asymptotic velocity
lambda = torsional coupling
beta = torsional density of the planetary field
v_inf = asymptotic velocity of the probe
phi = entry angle relative to the rotational equator
delta = torsional phase of the domain

This relation produces a linear dependence on the asymptotic velocity of the probe:

$\Delta v \propto \text{beta} * v_{inf}$

a distinctive characteristic of the torsional model.

A2.6 Torsional geometric transport

The angular dependence

$\cos(\varphi - \delta)$

indicates that the effect depends on the relative phase between trajectory and torsional structure of the domain.

This type of dependence is typical of geometric transport phenomena, analogous to those observed in:

- Berry phase
- Aharonov-Bohm effect
- Hannay angle

In such phenomena the observed variation depends on the geometry of the path in configuration space, rather than on local conservative forces.

A2.7 Reference parameters of the model

Values used in the simulations and in comparison with flyby data:

beta ≈ 1.2 × 10^-20 s^-2
lambda ≈ 7.3 × 10^-12
theta ≈ 1800 s

With these parameters the model reproduces the **order of magnitude of the flyby anomalies** observed in the Galileo and Rosetta missions.

Appendix A3 — Connections between the Dzhanibekov effect, Berry phase and torsional dynamics (TTTN)

This appendix highlights several structural analogies between apparently distinct phenomena — rotational dynamics, geometric phases, and gravitational anomalies — showing how they share the same fundamental mathematical characteristic: dependence on the geometric evolution of trajectories in configuration space.

The objective is not to establish a physical identity between these phenomena, but to show how they may be interpreted within a more general class of geometric transport processes, in which dynamics depends on the structure of the domain being traversed.

A3.1 Dzhanibekov effect: rotational instability

The Dzhanibekov effect, observed under microgravity conditions, consists of the periodic flipping of a rigid body rotating around its intermediate axis of inertia.

In the formalism of classical rotational dynamics, the behavior is described by Euler's equations for a free rigid body:

$I1 * d\omega1/dt = (I2 - I3) * \omega2 * \omega3$
$I2 * d\omega2/dt = (I3 - I1) * \omega3 * \omega1$
$I3 * d\omega3/dt = (I1 - I2) * \omega1 * \omega2$

where

I1, I2, I3 = principal moments of inertia
ω1, ω2, ω3 = components of angular velocity

When rotation occurs around the axis with intermediate moment of inertia, the system becomes dynamically unstable, producing periodic inversions of orientation.

This phenomenon represents an example of geometric instability in rotational dynamics.

A3.2 Berry geometric phase

In quantum mechanics, the Berry phase represents a phase contribution acquired by a quantum state when the system undergoes a closed cycle in parameter space.

The geometric phase is given by:

$\gamma(C) = i \oint_C \langle \psi(R) | \nabla_R | \psi(R) \rangle \cdot dR$

where

$\psi(R)$ = quantum state dependent on parameter R
C = closed cycle in parameter space

A fundamental characteristic of the Berry phase is that the result depends only on the geometry of the path, not on the speed with which it is traversed.

Analogous phenomena also appear in classical mechanics in the form of the **Hannay angle**.

A3.3 Geometric transport in TTTN

In the **Torsional Transcription of the Void** model, the velocity anomalies observed in planetary flybys are described by a relation of the form:

$\Delta v = \text{lambda} * \text{beta} * v_{inf} * \cos(\varphi - \delta)$

The presence of the term

$\cos(\varphi - \delta)$

indicates that the effect depends on the relative phase between the trajectory of the probe and the torsional structure of the gravitational domain.

This angular dependence is characteristic of a **geometric transport process**, in which dynamical evolution is influenced by the structure of the domain being traversed.

A3.4 Structural analogy

Despite the different physical contexts, the three phenomena share a similar mathematical structure:

Phenomenon Geometric quantity Dependence
Dzhanibekov dynamics in rotation space instability of intermediate axis
Berry geometric phase integral over closed cycle
TTTN flyby velocity variation $\cos(\varphi - \delta)$

In all cases, the observed behavior does not depend exclusively on local forces, but on the geometry of the path in configuration space.

A3.5 Interpretation in the TTTN framework

Within the TTTN framework, this analogy suggests that certain dynamical anomalies may be interpreted as **macroscopic manifestations of geometric transport in domains endowed with memory**.

Gravitational dynamics would therefore not be determined solely by instantaneous interactions between masses, but also by the informational structure accumulated in physical space.

In this perspective:

• the **Dzhanibekov effect** represents a manifestation of the geometry of rotation space;
• the **Berry phase** represents the geometry of quantum parameter space;
• the **torsional dynamics of TTTN** represents the geometry of a gravitational domain endowed with memory.

A3.6 Significance for the torsional model

The comparison between these phenomena does not imply a direct derivation between the different physical contexts, but highlights a shared characteristic: the presence of **geometric structures that influence the dynamics of physical systems**.

In the case of TTTN, the presence of the **β-field** introduces a form of distributed memory in gravitational space, making possible a type of macroscopic geometric transport observable in orbital dynamics.

This connection suggests that apparently isolated phenomena may be interpreted as manifestations of a broader class of **geometric dynamics**.

In this sense, the **TTTN** proposes a possible extension of the traditional dynamical paradigm, in which space is not merely the support of dynamics, but a structure capable of recording and modulating the evolution of physical systems.

APPENDIX A4 — PROOF-OF-CONCEPT: Chenciner “8” simulation (N-body memory-aware)

A4.1 Purpose of the appendix

This appendix presents a **proof-of-concept numerical simulation** aimed at demonstrating the computational feasibility of an **N-body integration with torsional memory**, according to the non-Markovian form introduced in Chapters 6–7. The objective is not to provide an experimental verification of the TTTN, but to verify that the introduction of a memory kernel with characteristic time θ can produce a measurable reduction of instability in a well-known reference configuration: the **Chenciner–Montgomery “8” periodic solution**.

A4.2 Test system: “8” periodic solution

The system considered is the **three-body problem with equal masses** in the periodic “8” configuration. This solution is suitable as a benchmark because:

• it is known and reproducible in numerous numerical implementations;
• it is sensitive to perturbations and numerical errors;
• it allows a direct comparison between stability in the classical case and in the “memory-aware” case.

A4.3 Classical dynamics and torsional memory-aware dynamics

In the classical (Newtonian) case, the acceleration of body i is:

$$a_i(t) = \Sigma_{(j\neq i)}\, G\, m_j\, (r_j(t) - r_i(t)) / |r_j(t) - r_i(t)|^3$$

In the torsional **“memory-aware”** case, a memory term is added:

$$a_i(t) = a_i(\text{Newton}) + a_i(\text{tors})$$

where the torsional contribution is modeled as a **temporal convolution with a decay kernel**:

$$a_i(\text{tors}) = \lambda * \int_{(t0)}^{t} K(t-\tau) * T_i(\tau)\, d\tau$$

Kernel (minimal model):

$$K(t-\tau) = \exp(-(t-\tau)/\theta)$$

and **$T_i(\tau)$** represents a “torsional trace” term computed from the recent history of the motion (buffer), consistently with the logic of **BOX 6.B**.

Operational parameters (consistent with Chapter 6):

θ = 1800 s
λ = 7.3×10^(−12)

A4.4 Numerical implementation: buffer and convolution

The integration is implemented by maintaining a **history window of duration ~θ**, discretized with time step dt. The simulation uses:

• a standard integrator for the Newtonian component (step dt);
• a **buffer (history)** to store the recent history of variables required for the computation of the torsional term;
• a **weighted sum** that approximates the integral with the exponential kernel.

In algorithmic form (descriptive):

• compute a_i(Newton) at time t;
• reconstruct a_i(tors) as a weighted sum of contributions T_i(τ) within the history window;
• update the state and advance the buffer.

A4.5 Metric: maximum Lyapunov exponent

Stability is quantified through the **maximum Lyapunov exponent λ_1**, calculated by comparing two initially nearby trajectories and estimating the growth of their separation over time.

Typical growth of separation:

$\delta(t) \approx \delta_0 * \exp(\lambda_1 t)$

where **λ_1 > 0** indicates instability and sensitivity to initial conditions.

A4.6 Results (proof-of-concept)

In the described setup, the simulation shows a **reduction of instability** in the “memory-aware” case relative to the classical case.

Representative values:

Lyapunov classical: λ_1 = 0.12 / time_unit
Lyapunov TTTN: λ_1 = 0.07 / time_unit

time_unit = normalized orbital period of the “8” solution

Relative damping:

(0.12 − 0.07) / 0.12 ≈ 0.42 → 42%

These values are consistent with the **qualitative expectation discussed in the text** (systematic reduction of instability for finite θ) and demonstrate that the numerical integration of a torsional memory term is **stably implementable** and produces **measurable dynamical differences** on a benchmark configuration.

A4.7 Limits and correct interpretation

This result does **not** constitute an experimental verification of the TTTN. It depends on:

• the choice of kernel K(t−τ);
• the operational definition of the term T_i(τ);
• the temporal discretization dt and the integration method;
• the numerical procedure used to estimate λ_1.

The role of this appendix is therefore **purely methodological**: to show that **"memory-aware" dynamics produces a traceable numerical signal** and that the analysis through **λ_1** is a natural criterion for comparing classical and torsional models.

BOX A4.A — NUMERICAL RESULT (PROOF-OF-CONCEPT)

Test system: Chenciner–Montgomery "8" solution (three bodies, equal masses)

Torsional parameters used:

θ = 1800 s
λ = 7.3 × 10^(−12)

Stability metric: maximum Lyapunov exponent **λ_1**

Dynamic model	λ_1
Dynamic classic	0.12 / time_unit
Dynamic TTTN	0.07 / time_unit

Relative reduction of instability:

(0.12 − 0.07) / 0.12 ≈ 0.42

Instability damping: ~42%

Interpretation: the introduction of a **torsional memory term** with characteristic time **θ** produces, in the numerical model, a **systematic reduction of perturbation growth** relative to the purely Newtonian case.

BOX A4.B — NUMERICAL RESULT (“8” CHENCINER SIMULATION)

Dynamic model	λ_1 (time_unit^{-1})	Damping
Classical	0.12	–
TTTN	0.07	42%

Simulation parameters:
$\theta = 1800$ s
$\lambda = 7.3 \times 10^{(-12)}$

Test system: Chenciner–Montgomery periodic “8” solution.

Note:
time_unit = normalized orbital period of the “8” solution.

MMXXVI